



Fertilizing capacity of cryopreserved sperm of *Chirostoma jordani* (Woolman, 1894)

Bustamante-González Jesús Dámaso^{1,*}, Gutiérrez-Díaz Dulce Leticia², Baca-Alejo Judith Sarai³, Figueroa-Lucero Gerardo¹, Arenas-Ríos Edith⁴, Hernández-Rubio María Cecilia⁵, Ávalos-Rodríguez Alejandro⁶

¹Planta Experimental de Producción Acuicola, Universidad Autónoma Metropolitana-Iztapalapa (UAM-I), CDMX 09340, México

²Maestría en Ciencias Agropecuarias, Universidad Autónoma Metropolitana-Xochimilco (UAM-X), CDMX 04960, México

³Escuela Superior de Medicina, Instituto Politécnico Nacional (IPN), CDMX 11340, México

⁴Laboratorio de Morfofisiología y Bioquímica del Espermatozoide, UAM-I, CDMX 09340, México

⁵Departamento de Zoología, Escuela Nacional de Ciencias Biológicas, IPN, CDMX 11340, México

⁶Laboratorio de Bioquímica de la Reproducción, UAM-X, CDMX 04960, México

Abstract

The genus *Chirostoma* is endemic from the Mesa Central of Mexico. It is conformed of 18 species and six subspecies. Five species are in some category of risk, because of this, *Chirostoma jordani* is an excellent model species to implement biotechnologies like gametes cryopreservation. Aim of present study was to evaluate fertilizing capacity of cryopreserved *C. jordani* sperm, as alternative to conservation and assisted reproduction in this specie and genus. Males and females were collected from wild Atlangatepec dam stock, Tlaxcala State, Mexico. Seminal quality was evaluated in fresh and cryopreserved semen with three cryoprotective agents (CPAs): 10% dimethyl sulfoxide (DMSO), 10% methanol (MeOH), 14% ethylene glycol (EG) and it was determined its post-thaw fertilizing capacity. Sperm motility percentage decreased during cryopreservation process ($p < 0.05$). There were not significant differences in post-thaw motility percentage between EG ($53.5 \pm 1.9\%$) and MeOH ($53.3 \pm 1.3\%$), but DMSO ($50.3 \pm 0.5\%$) was significantly different ($p < 0.05$). Results showed that 0.2 μL fresh semen were enough to fertilize 100% oocytes ($n = 60$). 10 μL DMSO and 5 μL MeOH and EG cryopreserved semen were necessary to fertilize oocytes 100% ($n = 60$) ($p < 0.05$). Cryopreservation and fertilization protocol for *C. jordani* sperm was efficient and it could be used for its assisted reproduction.

Keywords: Sperm cryopreservation, Cryoprotective agents, Fertilizing, Assisted reproduction

Introduction

The genus *Chirostoma* includes charales and white fish, they are endemic from the Mesa Central of Mexico; this one is composed of 18 species and 6 subspecies from Atherinopsidae,

controversial taxon in terms of its origin, colonization and diversification in the Altiplano Mexicano (Barbour, 1973). It is discontinuously distributed in lentic systems, along the Lerma River Basin (Hernández-Rubio et al., 2006). This basin is characterized by a high degree of urbanization and industrialization,

Received: Oct 20, 2023 Revised: Jan 11, 2024 Accepted: Jan 26, 2024

*Corresponding author: Bustamante-González Jesús Dámaso

Planta Experimental de Producción Acuicola, Universidad Autónoma Metropolitana-Iztapalapa (UAM-I), CDMX 09340, México

Tel: +52-56-10-93-52-43, E-mail: jesusbustamantegonzalez@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2024 The Korean Society of Fisheries and Aquatic Science

habitat fragmentation and exotic species introduction, that have decreased their fish populations size and others one are in some category of risk (SEMARNAT, 2019).

Chirostoma jordani (Woolman, 1894) has the major distribution of the *jordani* group (Barbour, 1973; Miller et al., 2005). It inhabits in clear and turbid waters from several lakes, rivers channels and dams, whit great load of sewage waters and xenobiotic contaminants that negatively affect biochemical and physiological functions of aquatic fauna, besides of insecticides, agrochemicals and fertilizers derived from agricultural activities (López-López et al., 2006).

C. jordani, because of its wide distribution and not be in any risk category, in agree NOM 059-SEMARNAT (2019), it can be used as model to implement conservation strategies, applicable of five species in risk of *Chirostoma* (*C. bartoni*, *C. charari*, *C. labarcae*, *C. promelas* and *C. riojai*). Sperm cryopreservation is inside these conservation strategies. Because of this technique allows to maintain cell viability and functionality at low temperatures, it has been recognized as an effective method for assisted reproduction in multiple fish species, either in captivity or in some risk category (Viveiros et al., 2015).

Fish semen cryopreservation involves mixing it in diluents that prevent motility and cryoprotective solutions, whit osmotic and nutritional properties, which together provide a suitable medium to minimize damage during freezing and thawing (Bustamante-González et al., 2019).

Currently, there is a specific protocol of sperm cryopreservation for the *C. jordani* species (Bustamante-González et al., 2021) where four cryoprotective agents (CPAs) effect was evaluated: dimethyl sulfoxide (DMSO), methanol (MeOH), ethylene glycol (EG) and glycerol (GL) in five concentrations: 2%, 6%, 10%, 14% and 16% v/v. Results showed that *C. jordani* sperm can be cryopreserved in MeOH 10% and EG14%, at 10 and 5 min equilibrium times whit $54.5 \pm 1.0\%$ and $53.1 \pm 1.0\%$ post-thaw motility percentages, respectively, thawed at 40°C . This demonstrates the progress and interest in implementing and improving the proposed cryopreservation protocols for the genus *Chirostoma*.

Therefore, aim of present investigation was to evaluate fertilizing capacity in cryopreserved sperm of *C. jordani*, as a viable alternative for its conservation and/or assisted reproduction and the genus.

Materials and Methods

Ethical statement

Present investigation was carried out under collection permission DGOPA. 07343.310810.4128 SEMARNAT, Mexico. The sample of organisms was reduced to the minimum necessary to cover the protocol.

Biological material collection

It was collected 600 sexually mature males (semen presence, weight 3.9 ± 0.4 g, 8.9 ± 0.3 cm total length) and 150 sexually mature females (ova presence, 4.5 ± 0.6 g, 8.5 ± 0.4 cm total length). Fish was collected in Atlangatepec dam, Tlaxcala, Mexico.

Fresh semen collection

Males were anesthetized with clove essence (0.01 mL/L water), before semen collection. Genital opening was cleaned with absorbent paper; semen was obtained by light pressure on abdominal and it was collected with a micropipette (10 μL) (Bustamante-González et al., 2018).

Sperm concentration estimation

Sperm concentration was determinate individually ($n = 100$), from a stock solution (94 μL NaCl 9%, 5 μL formaldehyde 4% plus 1 μL semen) (Bustamante-González et al., 2018). Sperm number was estimated (cells per/ μL) in Neubauer camera under microscope at 100X (Olympus Optical BX41TF®, Olympus, Tokyo, Japan).

Motility estimation

Sperm motility was determinate in each of 100 males. It was active 1 μL fresh semen in 15 μL dam water and it was observed under optical microscope at 100X (Bustamante-González et al., 2021). This volume allowed sperm activation and motility estimation. Sperm motility was evaluated by one person to standardize bias in measurements and estimated from 0% to 100% subjectively (Borges et al., 2005; Bustamante-González et al., 2018).

Sperm cryopreservation

Sperm cryopreservation was carried out according to the specific protocol of sperm cryopreservation for *C. jordani* proposed by Bustamante-González et al. (2021). This one consists of suspending 10 μL semen in an extender solution composed by 0.09 M sodium glutamate, 0.04 M fructose, 0.003 M magnesium

acetate and 0.05 M potassium acetate (310 mOsmol/kg) (Lake & Ravie, 1984) in 1:1 ratio, in each of DMSO 10%, MeOH 10% and EG 14%, whit 10, 10 and 5 min equilibrium times, respectively, at 4°C. Later, samples are transferred to 0.25 mL straws placed 3 cm above liquid nitrogen for 5 min. Straws are stored in liquid nitrogen (-196°C) in a thermos (Cryosystem series XC 20/20, Custom Biogenic Systems, Bruce Township, MI, USA). After 15 days, straws are thawed by immersion individually, in water at 40°C for 15 s. Samples are transferred to 0.1 mL microtubes, they are homogenized and it is determined motility percentage (1 µL sample plus 5 µL dam water, as activator solution) by triplicate, under Olympus Optical BX41TF® microscope at 100X.

Oocytes collection

Sexually mature females (n = 150), oocytes present, were anesthetized with clove essence (0.1 mL/L water) before oocytes extraction. Oocytes was obtained by light pressure on abdominal. First 60 oocytes were obtained from each female and they were distributed in Petri boxes (60 × 15 mm) by quadruplicate.

Fertilizing capacity in fresh

Fertilizing capacity in fresh was done as follow: 1) Eight groups were conformed, each with four Petri boxes and each box with 60 oocytes, 2) Fresh semen from 10 males pooled (90% motility percentage), 3) each group was fertilized with following semen volumes: 0.4, 0.2, 0.1, 0.08, 0.06, 0.04, 0.02 and 0.01 µL; each semen volume was activated with 2.0, 1.0, 0.5, 0.4, 0.3, 0.2, 0.1 and 0.05 µL dam water (Table 1), 4) after semen was activated, it was aspirated with a micropipette and poured directly onto the oocytes, in each Petri boxes, 5) fertilization (%) was determined

Table 1. Tested fresh semen volumes to fertilization in *Chirostoma jordani*

Fresh semen				
Group	Semen (µL)	Sperm concentration (cells per/µL)	Activating solution (µL)	Oocytes
1	0.4	720,000	2.0	60
2	0.2	360,000	1.0	60
3	0.1	180,000	0.5	60
4	0.08	144,000	0.4	60
5	0.06	108,000	0.3	60
6	0.04	72,000	0.2	60
7	0.02	36,000	0.1	60
8	0.01	18,000	0.05	60

by perivitelline space formation, after 10 minutes.

Fertilizing capacity post-thaw

Fertilizing capacity post-thaw was carried out as follows: 1) three groups were formed with four batches, each batch with four Petri boxes and each box with 60 oocytes, 2) 16 straws were post-thaw from each CPAs: DMSO, MeOH and EG, 3) motility percentage was verified (≥ 50%), 4) each batch was fertilized with different semen volume, in each group: batch 1: 10 µL, batch 2: 5 µL, batch 3: 2.5 µL and batch 4: 1.25 µL. Each semen volume was activated with 50, 25, 12.5 and 6.25 µL dam water (Table 2), 5) after semen was activated, it was aspirated with a micropipette and poured directly onto the oocytes, in each Petri boxes, 6) fertilization (%) was determined by perivitelline space formation, after 10 minutes.

Data analysis

Volume, sperm concentration and sperm motility were expressed as mean ± SD from fresh semen samples. Normal distribution data was verified by Shapiro-Wilk test ($p < 0.05$). One-way ANOVA was employed (95% confidence, $\alpha = 0.05$), to determine cryopreservation effect over sperm motility. Concentrations and CPAs effects over sperm motility percentages were analyzed through a one-way ANOVA (95% confidence, $\alpha = 0.05$) followed by Tukey test ($\alpha = 0.05$). Therefore, fertility (%) between fresh and cryopreserved semen volumes was analyzed (ANOVA, $p < 0.05$). Data were analyzed by SigmaPlot-14 program (Systat Software, San Jose, CA, USA).

Results

Fresh seminal quality

Average semen volume of *C. jordani* was 3.86 ± 1.53 µL, with a sperm concentration $1.8 \pm 0.3 \times 10^6$ µL⁻¹ and ≥ 95% motility percentage.

Sperm motility percentage in fresh and cryopreserved semen

Sperm motility percentage in fresh samples decreased during cryopreservation process ($p < 0.05$). There were not differences among EG ($53.5 \pm 1.9\%$) and MeOH ($53.3 \pm 1.3\%$) ($p > 0.05$) but DMSO varied significantly ($50.3 \pm 0.5\%$) (Fig. 1).

Sperm fertilizing capacity

A volume of 0.2 µL fresh semen was enough to fertilize 100% oocytes (n = 60) with a relationship 6,000:1 sperm: oocyte.

Table 2. Fertilizing capacity of cryopreserved *Chirostoma jordani* sperm

Cryopreserved semen						
Group	Batch	Volume (μL)	Sperm concentration per straw (cells per/μL)	Sperm motility per straw	Activating solution (μL)	Oocytes
DMSO	1	10.0	18,000,000	9,054,000	50.0	60
	2	5.0	9,000,000	4,527,000	25.0	60
	3	2.5	4,500,000	2,263,500	12.5	60
	4	1.25	2,250,000	1,131,750	6.25	60
MeOH	1	10.0	18,000,000	9,594,000	50.0	60
	2	5.0	9,000,000	4,797,000	25.0	60
	3	2.5	4,500,000	2,398,500	12.5	60
	4	1.25	2,250,000	1,199,250	6.25	60
EG	1	10.0	18,000,000	9,630,000	50.0	60
	2	5.0	9,000,000	4,815,000	25.0	60
	3	2.5	4,500,000	2,407,500	12.5	60
	4	1.25	2,250,000	1,203,750	6.25	60

DMSO, dimethyl sulfoxide; MeOH, methanol; EG, ethylene glycol.




Fig. 1. Mean ± SD fresh and cryopreserved sperm motility with DMSO, MeOH and EG. ANOVA, $F = 1,658.788$, $p < 0.001$; Tukey $p < 0.05$. Different letters show significant differences. DMSO, dimethyl sulfoxide; MeOH, methanol; EG, ethylene glycol.

Volumes higher were not different ($p > 0.05$) but fertilization success was less with minor volumes ($p < 0.05$) (Fig. 2).

Semen cryopreserve volumes were greater to fertilize same

number of oocytes ($n = 60$) ($p < 0.05$). It was needed 10 μL in samples cryopreserved in DMSO (150,900 sperm: oocyte (Fig. 3A), and 5 μL of MeOH and EG (79,950 and 80,250 sperm: oocyte, respectively) (Fig. 3B and 3C).

There was no significant difference ($p > 0.05$) among batches of oocytes in MeOH fertilized with 10 and 5 μL semen, but they were different from batches fertilized with 2.5 and 1.5 μL semen ($p < 0.05$) (Fig. 3B). Finally, samples oocytes cryopreserved in EG were not different fertilized with 10, 5 and 2.5 μL semen, but all of them were from batches fertilized with 1.25 μL semen (Fig. 3C).

Discussion

Cryopreservation had an effect over fertilizing capacity of semen, because their viability and motility decreased during thaw and post-thaw process, considerably. Because of this, it has been registered it is necessary a greater volume or concentration of semen to obtain percentages similar to those with fresh semen (Bozkurt & Yavaş, 2017; Ciereszko et al., 2014; Di Iorio et al., 2019; Du et al., 2018; Kommisrud et al., 2020).

In all cryopreservation protocol is a lost in motility percentage, due to sensitivity degree of sperm to CPAs toxicity, osmotic stress of extender solution over cell and oxidative stress of cryopreservation effect (Judycka et al., 2020; Nynca et al., 2019).

Osmotic stress is attributed to permeable CPAs (DMSO, MeOH, EG, GL, propanediol) are easily spread in cell (Gironi




Fig. 2. Mean ± SD fecundation regarding fresh semen volume. ANOVA, $F = 269.602$, Tukey $p < 0.05$. Different letters show significant differences.




Fig. 3. Mean ± SD fecundation regarding semen volume in each cryoprotective agents (CPAs). A: DMSO (ANOVA, $F = 270.375$, $p < 0.001$; Tukey $p < 0.05$), B = MeOH (ANOVA $F = 216.437$, $p < 0.001$; Tukey $p < 0.05$), C = EG (ANOVA $F = 264.648$, $p < 0.001$; Tukey $p < 0.05$). Different letters show significant differences. DMSO, dimethyl sulfoxide; MeOH, methanol; EG, ethylene glycol.

et al., 2020). Their principal function in cryopreservation is to prevent ice crystals formation inside cell. However, the CPAs can have next behavior during the process: 1) Ice is formed in extracellular space during cryopreservation. This gives rise to increase solutes concentration and causes an osmotic imbalance between inside and outside of cell. 2) Cooling rate should be slow enough to allow solutes and water are exchanged, in order to keep minimum imbalance osmotic. 3) When icy melts, extracellular space is hypotonic relative to inside cell, hence it can cause rapid ingress of water and give rise to cell lysis (Notman et al., 2006). Therefore, it is necessary to standardize equilibrium time according CPAs which depends on CPAs concentration and target species.

Besides, Gironi et al. (2020) and Kumari & Kashap (2019) report CPAs, in low concentrations, induces disorder of fatty acid tails of membrane phospholipids, in addition to increase phospholipid bilayer fluidity; high concentrations of these cause pores formation through membrane and their later increases cause plasma membrane disintegration. These results could support why sperm motility percentage depends of concentration and type CPAs. A previous study showed low concentrations of these agents induced lower motility percentage and there was a maximum motility value in higher CPAs, and from which it was observed a percentage decrease (Bustamante-González et al., 2021). This behavior may be due to high concentrations agents which form transient pores water or membrane disintegration, in extreme cases, both affect sperm viability (Gironi et al., 2020; Kumary & Kashap, 2019).

In addition, cryopreservation generates numerous structural and physiological changes to spermatozoa: thin membrane, increased fluidity, transient water pores through phospholipid bilayer, phospholipid bilayer breakdown, plasma membrane rupture, altered mitochondrial potential, reactive oxygen species production, DNA fragmentation, protein function, adenosine triphosphate (ATP) concentrations alterations, intracellular calcium homeostasis, among others, together affect fertilization (Ahn et al., 2018; Bustamante-González et al., 2019; Díaz et al., 2019; Figueroa et al., 2019; Gironi et al., 2020; Kumari & Kashyap, 2019; Lee-Estevez et al., 2019; Malcervelli et al., 2020; Schrader et al., 2016). Because of this, it is estimated 30% sperm motility loss for fertilization (Valipour, 2021). It was presented a loss about 50% in this study, similar to previous research (Bustamante-González et al., 2021); this motility loss may be due to cryopreservation protocol causes these structural and physiological changes to spermatozoa. These changes have been

evaluated in other fish species (Cabrita et al., 2014; Díaz et al., 2019; Figueroa et al., 2019; Sandoval-Vargas et al., 2020).

Success of a protocol cryopreservation is keeping sperm motility to fertilization. several researches have reported fertilizing capacity of sperm cryopreserved, mainly in model species of Salmonidae and Cyprinidae, where fertilizing capacity sperm decreases significantly during cryopreservation process, regarding oocytes fertilized with fresh semen.

Nevertheless, these researches have used same sperm concentration per oocyte, in fresh as cryopreserved, without considering that motility percentage, viability and ATP decrease with cryopreservation, significantly (Figueroa et al., 2019; Kommisrud et al., 2020; Sandoval-Vargas et al., 2020). For this reason, some authors recommend using excessive amounts of cryopreserved semen to increase fecundation (Tekin et al., 2007). Similar results were obtained in present study.

Sperm-oocyte ratio depends from species, mainly on assisted reproduction, since a greater number of oocytes can be fertilized with a lower sperm concentration. An advantage from assisted reproduction is to limit sperm displacement space so they invest their energy in penetrating the micropyle. This could explain differences in sperm number needed to fertilization among species, which is between 15×10^6 and 5×10^3 fresh sperm per oocyte, with a fecundation higher 90%. Similar results were obtained with fresh semen from *C. jordani*.

Species habitat is another important variable to consider. Different fertilization (%) have been obtained in brown trout populations that inhabit different rivers within same basin. This answer may explain sperm resistance to type and concentration of extender solution and CPAs (Ciereszko et al., 2014; Nynca et al., 2019), since both affect sperm motility. Results obtained in present research show that sperm concentration per oocyte depends on cryopreservation protocol, fertilization, species and even, among individuals of the same species, and they coincide with those obtained by other studies (Bozkurt & Yavaş, 2017; Ciereszko et al., 2014; Di lorio et al., 2019; Du et al., 2018; Kommisrud et al., 2020).

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was carried out with financial support was provided by Consejo Nacional de Humanidades, Ciencias y Tecnologías

(CONAHCYT) to the first author (BGJD, CVU: 566816) and is part of the project (CD.CBS.486.2021).

Acknowledgements

Present research is part of postdoctoral stay of first author (BGJD, CVU: 566816).

Availability of data and materials

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Ethics approval and consent to participate

This study conformed to the guidance of animal ethical treatment for the care and use of experimental animals.

ORCID

Bustamante-González Jesús Dámaso
<https://orcid.org/0000-0003-2912-5352>

References

- Ahn JY, Park JY, Lim HK. Effects of different diluents, cryoprotective agents, and freezing rates on sperm cryopreservation in *Epinephelus akaara*. *Cryobiology*. 2018;83:60-4.
- Barbour CD. A biogeographical history of *Chirostoma* (Pisces: Atherinidae): a species flock from the Mexican plateau. *Copeia*. 1973;3:533-56.
- Borges A, Siqueira DR, Jurinitz DF, Zanini R, Amaral F, Grillo ML, et al. Biochemical composition of seminal plasma and annual variations in semen characteristics of jundiá *Rhamdia quelen* (Quoy and Gaimard, Pimelodidae). *Fish Physiol Biochem*. 2005;31:45-53.
- Bozkurt Y, Yavaş I. Effect of different straw volumes and thawing rates on post-thaw quality and fertilization ability of cryopreserved common carp (*Cyprinus carpio*) sperm. *J Limnol Freshw Fish Res*. 2017;3:25-31.
- Bustamante-González JD, Cortés-García A, Ávalos Rodríguez A, Figueroa Lucero G, Arenas Ríos E, Gutiérrez MR. Semen evaluation in *Chirostoma jordani* (Woolman, 1894) and *Chirostoma humboldtianum* (Valenciennes, 1835), Mexican native species (Atheriniformes: Atherinopsidae). *Int J Aquatic Sci*. 2018;9:38-43.
- Bustamante-González JD, Rodríguez-Gutiérrez M, Cortés-García A, Arenas-Ríos E, Figueroa-Lucero G, Ávalos-Rodríguez A. Cryopreservation of *Chirostoma jordani* sperm, fish model for the conservation of the genus in Mexico. *J Appl Ichthyol*. 2021;37:278-84.
- Bustamante-González JD, Rodríguez-Gutiérrez M, Cortés-García A, Arenas-Ríos E, Figueroa-Lucero G, Ávalos-Rodríguez A. Fisiología y criopreservación del espermatozoide en teleósteos. *AquaTIC*. 2019;53:1-17.
- Cabrita E, Martínez-Páramo S, Gavaia PJ, Riesco MF, Valcarce DG, Sarasquete MP, et al. Factors enhancing fish sperm quality and emerging tools for sperm analysis. *Aquaculture*. 2014;432:389-401.
- Ciereszko A, Dietrich GJ, Nynca J, Dobosz S, Zalewski T. Cryopreservation of rainbow trout semen using a glucose-methanol extender. *Aquaculture*. 2014;420:275-81.
- Di Iorio M, Esposito S, Rusco G, Roncarati A, Miranda M, Gibertoni PP, et al. Semen cryopreservation for the Mediterranean brown trout of the Biferno River (Molise-Italy): comparative study on the effects of basic extenders and cryoprotectants. *Sci Rep*. 2019;9:9703.
- Díaz R, Lee-Estevez M, Quiñones J, Dumorné K, Short S, Ulloa-Rodríguez P, et al. Changes in Atlantic salmon (*Salmo salar*) sperm morphology and membrane lipid composition related to cold storage and cryopreservation. *Anim Reprod Sci*. 2019;204:50-9.
- Du C, Han YL, Shi YX, Zhu JQ. Cryopreservation of *Plagiognathops microlepis* sperm. *Cryobiology*. 2018;85:105-12.
- Figueroa E, Lee-Estevez M, Valdebenito I, Watanabe I, Oliveira RPS, Romero J, et al. Effects of cryopreservation on mitochondrial function and sperm quality in fish. *Aquaculture*. 2019;511:634190.
- Gironi B, Kahveci Z, McGill B, Lechner Bob Dan, Pagliara S, Metz J, et al. Effect of DMSO on the mechanical and structural properties of model and biological membranes. *Biophys J* 2020;119:274-86.
- Hernández-Rubio MC, Figueroa-Lucero G, Barriga-Sosa IDLA, Arredondo-Figueroa JL, Castro-Barrera T. Early development of the shortfin silverside *Chirostoma humboldtianum* (Valenciennes, 1835) (Atheriniformes: Atherinopsidae). *Aquaculture*. 2006;261:1440-6.
- Judycka S, Słowińska M, Nynca J, Liszewska E, Dobosz S, Ciereszko A. Oxidative stress in cryopreserved semen of sex-reversed female and normal male rainbow trout. *Aquaculture*. 2020;528:1-8.
- Kommisrud E, Myromslien FD, Stenseth EB, Zeremichael TT, Hofman N, Grevle I, et al. Viability, motility, ATP content and fertilizing potential of sperm from Atlantic salmon

- (*Salmo salar* L.) in milt stored before cryopreservation. *Theriogenology*. 2020;151:58-65.
- Kumari P, Kashyap HK. DMSO induced dehydration of heterogeneous lipid bilayers and its impact on their structures. *J Chem Phys*. 2019;151:215103-12.
- Lake PE, Ravie O. An exploration of cryoprotective compounds for fowl spermatozoa. *Br Poult Sci*. 1984;25:145-50.
- Lee-Estevez M, Herrera L, Díaz R, Beltrán J, Figueroa E, Dumorné K, et al. Effects of cryopreservation on cAMP-dependent protein kinase and AMP-activated protein kinase in Atlantic salmon (*Salmo salar*) spermatozoa: relation with post-thaw motility. *Anim Reprod Sci*. 2019;209:106133.
- López-López E, Sedeño-Díaz JE, Perozzi F. Lipid peroxidation and acetylcholinesterase activity as biomarkers in the Black sailfin Goodeis, *Girardinichthys viviparus* (Bustamante) exposed to water from Lake Xochimilco (Mexico). *Aquat Ecosyst Health Manag*. 2006;9:379-85.
- Malcervelli DM, Torres P, Suhevic JE, Cisale H, Fischman ML. Effect of different glycerol concentrations on phosphatidylserine translocation and mitochondrial membrane potential in chilled boar spermatozoa. *Cryobiology*. 2020;95:97-102.
- Miller RR, Minckley WL, Norris SN. *Freshwater fishes of México*. Chicago, IL: The University of Chicago Press; 2005.
- Notman R, Noro M, O'Malley B, Anwar J. Molecular basis for dimethyl sulfoxide (DMSO) action on lipid membranes. *J Am Chem Soc*. 2006;128:13982-3.
- Nynca J, Słowińska M, Judycka S, Ciereszko A. Maladaptation of trout spermatozoa to fresh water is related to oxidative stress and proteome changes. *Reproduction*. 2019;157:485-99.
- Sandoval-Vargas L, Dumorné K, Contreras P, Farias JG, Figueroa E, Risopatrón J, et al. Cryopreservation of coho salmon sperm (*Oncorhynchus kisutch*): effect on sperm function, oxidative stress and fertilizing capacity. *Aquaculture*. 2020;533:736151.
- Schrader AM, Cheng CY, Israelachvili JN, Han S. Communication: contrasting effects of glycerol and DMSO on lipid membrane surface hydration dynamics and forces. *J Chem Phys*. 2016;145:041101.
- Secretaría del Medio Ambiente y Recursos Naturales [SEMARNAT]. Norma Oficial Mexicana NOM-059-SEMARNAT-2019, Protección ambiental-Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en 2010 [Internet]. *Diario Oficial De La Federación*. 2019 [cited 2024 Feb 2]. https://www.dof.gob.mx/nota_detalle.php?codigo=5578808&fecha=14/11/2019#gsc.tab=0
- Tekin N, Seccer S, Akcay E, Bozkurt Y, Kayam S. Effects of glycerol additions on post-thaw fertility of frozen rainbow trout sperm, with an emphasis on interaction between extender and cryoprotectant. *J Appl Ichthyol*. 2007;23:60-3.
- Valipour J, Nashtaei MS, Khosravizadeh Z, Mahdavinezhad F, Nekoonam S, Esfandyari S, et al. Effect of sulforaphane on apoptosis, reactive oxygen species and lipids peroxidation of human sperm during cryopreservation. *Cryobiology*. 2021;99:122-30.
- Viveiros ATM, Nascimento AF, Leal AC, Gonçalves ACS, Ordão LH, Cosson J. Methyl glycol, methanol and DMSO effects on post-thaw motility, velocities, membrane integrity and mitochondrial function of *Brycon orbignyanus* and *Prochilodus lineatus* (Characiformes) sperm. *Fish Physiol Biochem*. 2015;41:193-201.