

Fish Aquat Sci. 2025;28(6):357-371 https://doi.org/10.47853/FAS.2025.e31 eISSN 2234-1757 Fisheries and Aquatic Sciences

Reproductive biology and feeding habits of the Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758) (Pisces, Cichlidae) in Lake Hawassa, Ethiopia

Teshome Belay^{1, 2, *}, Elias Dadebo¹, Girma Tilahun¹, Degsera Aemro³

¹ Department of Aquatic Sciences, Fisheries and Aquaculture, College of Natural and Computational Sciences, Hawassa University, Hawassa 05, Ethiopia ² Department of Animal Sciences, College of Agriculture and Natural Resource, Dilla University, Dilla 419, Ethiopia

³ Department of Fisheries and Aquatic Sciences, School of Fisheries and Wildlife, College of Agriculture and Environmental Sciences, Bahir Dar University, Bahir Dar 5501, Ethiopia

Abstract

The study conducted from February 2023 to January 2024 aimed to investigate the reproductive biology and feeding habits of Nile tilapia Oreochromis niloticus in Lake Hawassa, Ethiopia. 816 mixed sexes of O. niloticus were considered for reproductive biology, sampling involved gillnets with mesh sizes of 6, 8, 10, and 12 cm and random samples from landing sites. To get different size ranges additional data were collected for feeding habits and morphometric relationships using smaller sizes of gillnets (2 and 2.5 cm), hand nets (6 mm size), and randomly from local fishers gathered with hooks, resulting in a total sample of 987 collectively. Microsoft Excel, SPSSv27, and MATLAB software were used for data analysis. The results showed that the sex ratio (1:0.93) did not showed deviate from the theoretical 1:1 ratio (χ^2 , p > 0.05). The TL_{50} % maturity was 18.5 cm for males and 19.1 cm for females. O. niloticus from Lake Hawassa showed year-round breeding with peaking in January and February. Fecundity ranged from 105 to 1,541 eggs/ripe fish. Phytoplankton was the most frequent (95.72%), contributed (88.2%), and preponed (96.6%) in the stomach of O. niloticus. Ontogenetic dietary shifts were not statistically significant (C > 0.6). Significant dietary variations were noted between rainy and dry months, particularly for the occurrence of macrophytes (χ^2 , p < 0.05) and volumetric contributions of detritus (t-test, p < 0.05). O. niloticus from Lake Hawassa showed positive allometric growth (b > 3) and was significantly different from 3 (t-test, p < 0.001). The mean relative condition factors showed significant variation by sampling month (ANOVA, p < 0.05). The result concluded that O. niloticus from Lake Hawassa is categorized as a phytoplanktivorous trophic category. Based on the present findings, it is recommended to stop the harvesting of O. niloticus below 19.1 cm and enforce fishing closures during peak breeding months (January and February). The insignificant dietary shift and phytoplankton dominance in the diet of O. niloticus suggest that a parallel study on phytoplankton biomass and primary production, as a proxy of yield and management of O. niloticus stock in Lake Hawassa, should be considered for future studies.

Keywords: Feeding habits, Lake Hawassa, Oreochromis niloticus, Phytoplanktivorous, Reproductive biology

Received: Oct 6, 2024 Revised: Dec 8, 2024 Accepted: Dec 30, 2024 *Corresponding author: Teshome Belay

Department of Aquatic Sciences, Fisheries and Aquaculture, College of Natural and Computational Sciences, Hawassa University, Hawassa 05, Ethiopia Tel: +251-934967894, E-mail: teshimeansc@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (http://creativecommons.org/licenses/bync/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited. Copyright © 2025 The Korean Society of Fisheries and Aquatic Science

Introduction

The Oreochromis niloticus fish species is the most preferred for consumption in Ethiopia, making up 49% of the total captured fish production (Tesfaye & Wolff, 2014). In Lake Hawassa, 90% of the total fish production is attributed to O. niloticus (Reyntjens & Wudneh, 1998). Due to increasing fishing pressure, overfishing of O. niloticus stock from Lake Hawassa has been continuously identified (Tekle-Giorgis, 2018). The ideal fishing effort for O. niloticus was determined to be 700 gill nets/day (Reyntjens & Wudneh, 1998). But overfishing was evident, with 1,954 gill nets/day found in the Lake, more than twice the recommended effort (Tekle-Giorgis et al., 2017). It was seen that 22.7% of O. niloticus stock from Lake Hawassa was harvested before attaining the length at first maturity (TL_{50}) (Muluye et al., 2016). To overcome the threats, another researcher advised that the fishing effort be reduced to 650 gillnets/day to obtain a safe and optimal bio-economical yield of O. niloticus from the lake (Tekle-Giorgis, 2018). Additionally, aggressive fishing with mesh sizes smaller than 7.5 cm is jeopardizing the harvest of small-sized fish species from the lake (Dadebo et al., 2023). Despite these concerns, no detailed study has assessed the current status of O. niloticus' reproductive biology in Lake Hawassa. Thus, a comprehensive analysis of the reproductive biology is essential to understand the current status of O. niloticus in the lake.

In terms of feeding habits, studies indicate that O. niloticus from Lake Hawassa is an herbivorous trophic category, concentrating on adult sizes ranging from 18-32 cm (Getachew & Fernando, 1989). Juvenile O. niloticus from Lake Hawassa exhibit a preference for zooplankton and some zoobenthos (Tudorancea et al., 1988). However, there is a notable gap in research regarding the dietary overlap between juvenile and adult O. niloticus, hindering the understanding of intraspecific feed resource competition. Therefore, concurrent studies of juvenile and adult O. niloticus are imperative to comprehend intraspecific competitions and ontogenetic dietary shifts in O. niloticus from Lake Hawassa. Furthermore, Lake Hawassa has suffered from pollution-related effects (Tilahun, 2023), leading to a decline in zooplankton abundance and biomass (Beyene et al., 2022), including important species such as Rotifers, Cladocera's, and Cycloid copepods, previously identified as a dietary source for O. niloticus juveniles (Tudorancea et al., 1988).

There are research reports that certain fish species from

Lake Hawassa exhibit a discernible shift in trophic behavior over time, necessitating further investigation into the dietary patterns and feeding habits of O. niloticus. Notably, initial perceptions regarding the carnivorous nature of larger fish species such as the African catfish Clarias gariepinus (Burchell, 1822) indicated insignificant reliance on plankton (Dadebo, 2000). Subsequently, C. gariepinus was reclassified as an omnivorous species, with significant contributions from phytoplankton and zooplankton (Tekle-Giorgis et al., 2016). This transition in feeding behavior underscores the need to reassess the current dietary habits of O. niloticus in the lake. In addition to the previously published information, the first study concerning the levels of microplastic concentration in O. niloticus muscle and gastrointestinal tract was identified (Demsie & Yimer, 2024). All these pollution-related concerns highlight the importance of researching the growth (b) of O. niloticus in Lake Hawassa. Consequently, the current study is designed to elucidate the reproductive biology and feeding habits of O. niloticus from Lake Hawassa. The resultant findings will prove invaluable for the development of stock assessment and modeling, population dynamic studies, Ecopath modeling, and the provision of updat ed information for management decisions.

Materials and Methods

Descriptions of the study area

Lake Hawassa is located within the Central Rift Valley Lakes of Ethiopia at 275 kilometers away from Ethiopia's capital city, Addis Ababa, in latitudes 6°49'-7°15'N and longitudes 38°17'-38°44'E (Dadebo et al., 2023). Though River Tikurwuha feeds the lake, it lacks a distinct outflow (Admassu, 1996). When the impact of wind is taken into consideration over the dry months, the lake water exhibits a high level of nutrient mixing (Tilahun & Ahlgren, 2010). The lake is characterized as a heaven of hundreds of phytoplankton species (Kebede & Belay, 1994), and 22 known zooplankton species dominated by rotifers (Beyene et al., 2022). Six distinct fish species are available in the lake (Dadebo, 2000). Three fish species are important for commerce: O. niloticus, C. gariepinus, and African big barb Labeobarbus intermedius (Ruppell, 1836). The remaining three species, the straightfine barb Enteromious paludinosis (Peters, 1852), stone lamping minnow Garra quadrimaculata (Ruppel, 1835), and black lampeye Aplocheilichthys antinorii (Vinciguerra, 1883), are ecologically significant and contribute as a food item for large fish species like C. gariepinus (Dadebo, 2000; Fig. 1). Due



Fig. 1. Geographic locations of Lake Hawassa. Adapted from Arc Gis Software with CC-BY-NC (Map generated by the author using Geographic Information System (GIS) software).

to the lake's exposure to several human activities, both the point and non-point pollutant sources are regarded as a threat to aquatic biota (Tilahun, 2023).

Fish sampling

Multiple approaches for collecting data have been applied to achieve the stated objectives. The gill net was one of the sampling techniques. The gill net was set at 5:00 PM and retrieved at 7:00 AM following a 14-hour stay. Gillnets with 6, 8, 10, and 12 cm stretched mesh sizes, with 50 × 1.5 m, were used for the reproductive biology study. A randomized sampling was also taken from the two major landing sites, Amoragedel and Tikurwuha. To get a wide size range for understanding the species' feeding habit and morphometric relationship, additional gillnets with stretched sizes of 2.0 and 2.5 cm were used. Further, hand-net-based sample collection techniques were also employed to get a smaller size of *O. niloticus* from the shoreline area. A random sample from local fishers who collect by hook was also collected.

Morphometric measurement

Using a portable weighing balance and measuring board, the weight and length of the samples were measured to the nearest 1.0 g and 0.1 cm, respectively. The fish specimens were dissected; the gonads and stomach were removed and preserved in a 5% formaldehyde solution. On the other hand, smaller-sized *O. niloticus*, measuring less than 10 cm, were measured

and dissected at the Hawassa University Fisheries Laboratory using a 0.1g scale sensitive balance and a 0.1 cm measuring board for diet composition analysis. The preserved gonads and stomach samples were taken directly to the fisheries laboratory at Hawassa University for additional examination of their reproductive and feeding biology. Female ripe ovaries were given a specific treatment in the lab. A surgical blade with a 22 mm diameter was used to dissect the ovary longitudinally and preserved until further procedures were applied.

Reproductive biology analysis

Gonad stage identification: The maturity stage of each sample was grouped into five points of maturity (Holden & Raitt, 1974). Finally, the data were expressed in percentages after categorizing them as non-breeding (I and II) and breeding (III, IV, and V).

$$MS_i \% = \frac{MS_i}{\sum_{i=1}^5 MS_i} \times 100 \tag{1}$$

where, MS_i %= the percent of *O. niloticus* at maturity sge *i*, MS_i = the number of fishes at maturity stage *i*, and ΣMS_i , is the total number of *O. niloticus* from maturity stage I to V.

Sex ratio: The sex ratio was calculated after categorizing the data into sampling months, size class, and total population. Finally, the sex ratio was compared using the chi-square test (Tadesse, 1997).

$$\chi^2 = \sum \frac{(Oi - Ei)^2}{Ei}$$
(2)

 x^2 is the chi-square value, O_i is the observed value, and is the expected value.

Length at 50% maturity: The length at 50% sexual maturity (TL_{50}) was calculated based on the logistic regression model (King, 2013). The numbers of mature and immature O. niloticus were identified after grouping into 2 size classes. The modal class (mid-length) was plotted as an X variable. Whereas the natural logarithm of the proportion was represented as a "Y" variable, which was calculated as $_{LN}\left|\left(\frac{1}{p}\right)^{-1}\right|$, where LN represents the natural logarithm and p represents the proportions of matured fish per total sample in each size class. After identifying the value of X (mid-length) and Y (LN transformed proportion), X-Y scatterplots were run using the Excel 10 sheet. The slope (b) and intercept (a) were extracted from the X-Y scatter plot graph which represents the mid-length and the proportions respectively. After this step, the specific value of TL_{50} was calculated using the equation $TL_{50} = \left[\left(\frac{-b}{a} \right) \right]$, where *b* represents slope and a represents intercept. To get the probability of being mature in each modal class, the equation was run as

$$Pm = \frac{100}{\left(1/_{EXp} \left(a - (b * X)\right)\right)}$$
(3)

where pm is the probability of being mature. Finally, the graph of the sigmoid curve was set with MATLAB software (MathWorks, Natick, MA, USA).

Breeding season: The ripe ovaries were extracted from the total sample for each sampling month to observe the specific frequency of ripe fish. The frequency of males and females of *O. niloticus* with ripe ovaries was plotted against the sampling month in percentage to show their peak and non-peak breeding seasons (Admassu, 1996).

Fecundity: For estimating the fecundity potentials, only ripe females (gonad stage V) were considered. The longitudinally preserved samples of each ovary were gently shaken and washed with tap water continuously to remove the ovarian membrane. Finally, total counting was applied for each egg from each ovary to know its fecundity potential.

Feeding habit analysis

Diet composition analysis: The preserved prey items were transferred with a known volume of a graduated cylinder test tube. Once the overall volume was known, the samples were shaken to homogenize the stomach contents. Larger specimens were identified using a dissecting microscope (LEICA MS), and smaller prey items were identified using a compound microscope (LEICA DME) with a magnification power of 100X. Finally, the results were interpreted based on the frequency of occurrences, volumetric contributions, and index of preponderances.

Frequency of occurrence: The number of specific preys per non-empty stomach was counted to understand how much it occurred in the total non-empty sample. The occurrences of each prey were expressed as a percentage (Hyslop, 1980).

$$\%FO = \left(\frac{TNS_i}{TNSP_i}\right) \times 100\tag{4}$$

where, FO = frequency of occurrences of specific prey from non-empty *O. niloticus* stomach, $TNSP_i$ = total number of *O. niloticus* stomach with specific prey item *i*, and TNS_i = total number of nun-empty *O. niloticus* considered for feeding habit analysis.

Volumetric contribution: The volumetric contributions of each food item per total volume of the non-empty stomach sample were expressed as percentages (Bowen, 1996).

$$V_i\% = \frac{VSP_iFS}{VF_iFS} \times 100$$
(5)

where V_i % indicates the volumetric contribution of prey item *i* in percentage, VSP_iFS indicates the volume of a specific prey item found in all specimens, and VF_iFS indicates the volume of all food items found in all non-empty stomachs.

Index of preponderance: The index of preponderance was calculated based on the frequency of occurrences and volumetric contribution. It was expressed as a percentage (Natarajan & Jhingran, 1961).

$$IoP\% = \frac{V_i x F_i}{\sum (V_i F_i)} \times 100$$
(6)

where F_i is the frequency of occurrence of food item *i*;*IoP*% is the index of preponderance of food item *i* in percentage and V_i is the percentage composition by volume of food item; *i*;*n* is the number of prey types, to facilitate comparisons among species.

Seasonal variations of diet compositions: The diet compositions were once again grouped and categorized based on rainy and dry months. Frequency of occurrences, volumetric contributions, and index of preponderance methods were used to compare the diet composition variations among seasons. Furthermore, the major prey item was statistically crosschecked using a chi-square test for the frequency of occurrences and an independent sample *t*-test for volumetric contributions, respectively (Dadebo et al., 2014).

Ontogenetic dietary shift and dietary overlap analysis: To determine the size-based ontogenetic dietary shift, the samples were grouped into 4 different size classes depending on their total length, and the percentage of volumetric contribution was calculated. For the first group, samples < 10 cm, the second group ranged from 10–19.9 cm, the third group was 20–29.9 cm, and the fourth group was considered above 29.9 cm. The volumetric contributions of major prey items were calculated in each size class, plotted with a stacked chart to indicate their contributions, and expressed as percentages. The dietary overlap was also considered to crosscheck the intra- and interspecific competitions. For the present study, intraspecific competitions based on size class comparisons were cross-checked depending on the major prey items contributed for each size class. The comparisons of dietary overlap of major prey items were compared between < 10 cm and 10–19.9 cm, < 10 cm and 20– 29.9 cm, < 10 cm and \ge 29.9 cm, 10–19.9 cm and 20–29.9 cm, $10-19.9 \text{ cm} \text{ and} \ge 29.9 \text{ cm}, 20-29.9 \text{ cm}, \text{ and} \ge 29.9 \text{ cm} \text{ size class}$ were considered. Mathematically, dietary overlap was calculated based on the Shooner dietary overlap index (Schoener, 1970).

$$C = 1 - \frac{1}{2} \left(\sum \left| p_{xi} - p_{yi} \right| \right) \tag{7}$$

where *C* is Schoener's similarity index metric, P_{xi} and P_{xy} and are the proportions of diet item *i* in the stomach of species *x* and *y*, respectively.

Length-weight and length-length relationships

For length-weight relationship analysis, the data was sorted and run using the power function after removing the outliers as follows: $TW = aTL^b$ (Pauly, 1984), where TW= total weight of fish (g), TL = total length of fish (cm), a = the intercept of the regression line, and b= the exponent (slope) of the regression line. Based on the value of the exponent (*b*), the fish growth is categorized into three groups (Pauly, 1984). When the slope value b = 3, the species has isometric growth; b < 3, the species follows a negative allometric growth; and b > 3, positive allometric growth. Considering their length-length relationship, linear regression comparisons were applied (Zar, 1999). The value was set as Y = aX + b, where Y and X represent the standard length (SL) and TL of *O. niloticus*, a is the intercept, and b is considered as the slope of a regression.

Relative condition factor

The interpretations of the *O. niloticus* health condition depend on the exponent (*b*) value generated from the length-weight relationship. Fulton's condition is considered for isometric growth (b = 3), whereas, for non-isometric growth (b \neq 3), and statistically significant difference from 3, the relative condition factor is considered as a benchmark for understanding the health of fish species (Jones et al., 1999). The comparisons were run both within the sampling month and sex-based comparisons as indicated below.

$$RCF = \left(\sqrt[w]{aTL^b} \right), (Zar, 1999), \tag{8}$$

where RCF indicates the relative condition factor, w represents total body weight, and a, and b describe constants and slopes of the length-weight relationship, respectively.

Data summarization, analysis, and presentation

The data was analyzed with descriptive statistics using IBM SPSSv27 (SPSS, Chicago, IL, USA) and MS-Excel 10 (Microsoft, Redmond, WA, USA). For sex ratio comparison, the chi-square test was applied. The exponents (b) of the length-weight relationship were taken from the logarithmic transformed data and tested using a student's *t*-test with a 95% confidence interval to know whether the value is significantly different from the expected cube value (3). The equation used for *t*-test comparisons of the exponent (b) is calculated as:

$$ts = ((b-3) / sd_b), (Zar, 1999),$$
 (9)

where ts is the *t*-calculated value, b is the slope, and sd_b is the SE, obtained from the log-transformed data's regression analysis. MATLAB software (MathWorks) was used to draw the graph of TL_{50} . Regarding the diet compositions, descriptive statistics using Microsoft Excel 10 (Microsoft) were used. The chi-square test for frequency of occurrences and independent sample *t*-test for volumetric contributions were applied to test whether the diet composition shows a significant variation between rainy and dry months. The dietary overlap was tested using Schoener's dietary overlap index to compare the intraspecific competition. The index was concluded as strong dietary overlap if the value is above 0.6 and considered as biological significance.

Result

Reproductive biology of Oreochromis niloticus

Sex ratio: From the total 816 samples collected for reproductive investigation, about 386 (47.3%) were females, and 430 (52.7%) were males. The result revealed that more males than females were caught. Nevertheless, comparisons of their sex ratio (1:0.93) were not statistically significant from the hypothetical 1:1 ratio (χ^2 , p > 0.05; Table 1). However, based on size-based comparisons, the result indicates that the sex ratio demonstrates a significant difference in a size range of 25.8 to 27.7 cm and from 27.8 to 30.7 cm (χ^2 , p < 0.05; Table 2).

Stage of maturity and length at first maturity: from the total sample size, 78.43% of *O. niloticus* was mature (breeding), whereas 21.57% was immature (non-breeding). Males' *O. niloticus* revealed that 23.95% were immature (non-breeding) and 76.05% were mature (breeding). Of the females' *O. niloticus*, 18.65% and 78.53% were immature and mature, respectively. The findings showed that in Lake Hawassa, more immature male *O. niloticus* were caught than female *O. niloticus*. Mature male *O. niloticus* was recorded at 17.8 cm in length and 123 gm in weight. Whereas female *O. niloticus*, was starting to mature at 17.4 cm in length and 103 gm in weight, respectively. In the current investigation, the TL_{50} % maturity for male and female *O. niloticus* was 18.5 cm and 19.1 cm, respectively. The result indicates male *O. niloticus* start to mature faster compared to females (Fig.2).

 Table 1. Sex ratio of Oreochromis niloticus at different

 sampling months (n = 816) rom Lake Hawassa

Months	Male	Female	Total	Sex ratio	χ²
February	59	53	112	1:0.90	0.32
March	27	31	58	1:1.15	0.28
April	44	31	75	1:0.83	2.25
May	30	44	74	1:1.47	2.65
June	30	30	60	1:1.00	0.00
July	27	26	53	1:0.96	0.02
August	33	20	53	1:0.61	3.19
September	32	25	57	1:0.78	0.86
October	39	35	74	1:0.90	0.22
November	37	26	63	1:0.70	1.92
December	41	29	70	1:0.88	2.06
January	31	36	67	1:1.16	0.37
Total	430	386	816	1:0.93	2.37

Table 2.	Sex ratio of	Oreochromis	niloticus a	t different	size
class (n =	= 816) from L	ake Hawassa			

Size class	Male	Female	Total	Sex ratio	χ²
13.8–15.7	1	1	2	1:1	0.00
15.8–17.7	1	2	3	1:2	0.33
17.8–19.7	12	6	18	1:0.5	2.00
19.8–21.7	58	75	133	1:1.29	2.17
22.8–23.7	170	154	324	1:0.91	0.79
23.8–25.7	138	115	253	1:0.83	2.09
25.8–27.7	29	15	44	1:0.52	4.45*
27.8–30.7	17	6	23	1:0.35	5.26*
30.8–32.7	3	8	11	1:2.67	2.27
≥ 32.8	1	4	5	1:4.00	1.80
Total	430	386	816	1:0.93	2.37

* Values are indicates the compressions show a significant difference (p < 0.05).



Fig. 2. Length at first maturity of *Oreochromis niloticus* from Lake Hawassa.

Breeding season: In Lake Hawassa during the present study, the *O. niloticus* breeding month indicates the species has the potential to breed throughout the year in both sexes. In both sexes of *O. niloticus*, the peak breeding seasons extended from February to April (Fig. 3). The peak breeding seasons for Females' *O. niloticus* were seen by January (20.0%) and February (21.9%). February (19.9%), April (11.8%), and January (11.2%) respectively were also the peak breeding seasons of males' *O. niloticus*.

Fecundity: The smallest length of a female *O. niloticus* with a ripe gonad was 17.4 cm and 102 g, respectively. The fecundity potential ranged from a minimum value of 105 eggs/ripe fish



Fig. 3. Breeding season of *Oreochromis niloticus* as indicated by the percentage of fish with ripe gonads (a), rainfall (bar chart b), maximum (●), and minimum (○) temperature of Hawassa surrounding.

to a highest value of 1,541 eggs/ripe fish, with a mean fecundity of 556 eggs/ripe fish. Based on the regression analysis, there is a significant interaction (ANOVA, p < 0.001) and a linear association between gonad weight and fecundity. The correlation analysis also showed a strong interaction between gonad weight and fecundity (r = 0.881). The equation was F = 57.635GW+105.88, n = 108, where F indicates fecundity, and GW indicates the gonad weight of *O. niloticus*.

Food and Feeding habits of *Oreochromis niloticus* from Lake Hawassa

Diet composition: From the 987 total samples collected for food and feeding habit analysis, 28 samples (2.84%) were empty, and the remaining 959 specimens (97.16%) stomachs were nonempty, which consists of several food items. Seven major food items were identified: phytoplankton, detritus, macrophytes, zooplankton, nematodes, fish eggs, and digested muscle (Table 3). Phytoplankton occurred to 95.72%; volumetrically,

Table 3. Frequency of occurrences, volumetric contributions, and index of the preponderance of various food items in the
diet of <i>Oreochromis niloticus</i> from Lake Hawassa (n = 959)

	Frequency of	Frequency of occurrences (FO)		Volumetric contribution (Vol)		Index of preponderance (IP)	
Food Item	FO	%FO	Vol	%Vol	IP	%IP	
Phytoplankton's	918*	95.72*	3,284.43*	88.20*	8,442.79*	96.60*	
Diatom	761	79.35	781.95	21.00	1,666.27	19.06	
Green Algae	618	64.44	678.43	18.22	1,174.02	13.43	
Bluegreen Algae	796	83.00	1,665.75	44.73	3,712.83	42.48	
Dinoflagellates	216	22.52	85.69	2.30	51.83	0.59	
Euglenoids	729	76.02	72.32	1.94	147.63	1.69	
Zooplankton's	152*	15.85*	62.51	1.68*	26.61*	0.30*	
Cladocerans	68	7.09	19.04	0.52	3.69	0.04	
Rotifers	62	6.47	25.73	0.69	4.47	0.05	
Copepods	42	4.38	17.74	0.48	2.09	0.02	
Detritus	298*	31.07*	266.53*	7.16*	222.40*	2.54*	
Macrophyte	182*	18.98*	94.23*	2.53*	48.02*	0.55*	
Nematode	6*	0.63*	1.60*	0.04*	0.03*	0.00*	
Fish Egg	4*	0.42*	11.75*	0.32*	0.13*	0.00*	
Digested Muscle	3*	0.31*	2.86*	0.08*	0.02*	0.00*	
Total				100.00	8,740.003	100.00	

* Values are mean represents major food item.

contributed to 88.20%, and the index of preponderance also shows 96.60% compared to the total food items. Compared to different types of phytoplankton, the non-filamentous types like blue-green algae and diatoms were the most preferred food item by *O. niloticus*. Detritus and macrophytes are also the second and third most preferred food items occurrence in 31.07% and 18.98%, with volumetric contributions to 7.16% and 2.53%, and index of preponderance 2.54% and 0.55%, respectively. Zooplankton is also the intermediate preferred food item, which occurred in 15.85%, volumetrically contributed to 1.68%, and the index of preponderance recorded 0.30%. However, nematodes, fish eggs, and digested muscles were the least contributors compared to the other preferred food items.

Ontogenetic dietary shift and dietary overlap: The diet composition of *O. niloticus* from Lake Hawassa indicated a slight ontogenetic dietary shift throughout their lifetime (Fig. 4). However, those slight shifts of feed resources have no significant change in all size classes which showed a strong dietary overlap (C > 0.6; Table 4). The contributions of phytoplankton increase when their size class increases. At smaller sizes below 10.0 cm, phytoplankton contributed to 62.63%; whereas in the size



Fig. 4. Ontogenetic dietary shift *Oreochromis niloticus* from Lake Hawassa.

Table 4. Dietary overlap between different size classes	of
Oreochromis niloticus from Lake Hawassa	

Size class (cm)	Shooner dietary overlap index (C)
< 10 and 10–19.9	0.81
< 10 and 20–29.9	0.73
< 10 and > 29.9	0.65
10–19.9 and 20–29.9	0.88
10–19.9 and > 29.9	0.79
20–29.9 and > 29.9	0.91

class10.0–19.9 cm, the phytoplankton contributed to 77.32%; at 20.0–29.9 cm, it contributed to 88.90%; and the larger size above 30.0 cm contributed to 98.05%. In a contradictory way, the contributions of zooplanktons at smaller sizes below 10.0 cm contributed more, and their contributions slightly decreased when their sizes increased. On larger size classes of *O. niloticus* above 30.0 cm, zooplanktons have no contributions, and their size class increases.

Seasonal variations: O. niloticus from Lake Hawassa showed slight diet variations in rainy and dry seasons. During the rainy season, the species preferred phytoplankton > detritus > macrophyte > zooplankton > nematode > fish egg > digested muscles (Table 5). However, except for macrophytes, prey occurrences in the rainy season didn't significantly differ from the dry season (χ^2 , p > 0.05). Similarly, except for detritus, the volumetric contributions of prey items didn't show a significant difference from the dry season contributions (*t*-test, p > 0.05). In the rainy season, phytoplankton occurred in 96.20%, volumetrically contributing to 87.70% and 96.36% on the index of preponderance. Blue-green algae were the major contributing prey item, compared to other phytoplankton taxa. Detritus was the second most preferred prey item in the rainy season, which recorded 32.07%, 8.25%, and 3.02% on the frequency of occurrences, volumetric contributions, and index of preponderance, respectively. Macrophyte and zooplanktons are intermediate contributors, which represent 15.33%, 1.91%, 0.33%; and 15.05%, 1.65%, and 0.82% in frequency of occurrences, volumetric contributions, and index of preponderance, respectively. Nematodes, fish eggs, and digested muscles are the least contributors compared to the major food items in the rainy season.

During the dry season, the contributions of diet com-positions indicate phytoplankton > macrophytes > detritus > zooplankton > fish eggs. Unlike the rainy season, during the dry season, macrophyte is considered the second most important food item, contributing to 29.44%, 4.20%, and 1.42% with the frequency of occurrences, volumetric contributions, and index of preponderance respectively. Detritus is also an intermediate contributor during dry season, which occurred in 28.23% and accounted for 4.18%, and 1.35% on the index of preponderance. Regarding the zooplankton, their occurrence (18.15%), contributions (1.77%), and preponderance (0.37%) showed a slight increase in the dry season compared to the rainy season. Unlike the rainy season, fish eggs contribute a very small amount, and there were no contributions from nematodes and digested muscles in dry seasons.

Food items	Frequency of occurrences (%)		Volumetric contribution (%)		Index of preponderance (%)	
	Rainy	Dry	Rainy	Dry	Rainy	Dry
Phytoplankton's	96.20ª	94.35°	87.70a	89.55ª	96.36 [*]	96.86*
Diatom	77.07	85.89	20.31	22.84	17.88	22.48
Green algae	63.43	67.34	19.26	15.43	13.96	11.91
Blue-green algae	84.25	79.44	43.44	48.19	41.79	43.89
Dinoflagellates	25.60**	13.71**	2.72**	1.17**	0.80**	0.18**
Euglenoids	16.88**	20.56**	1.95**	1.92**	0.38**	0.45**
Zooplanktons	15.05ª	18.15°	1.65°	1.77ª	0.28*	0.37*
Cladocera	6.19	9.68	0.48	0.61	0.03	0.07
Rotifers	6.47	6.45	0.64	0.99	0.05	0.04
Copepods	4.36	4.44	0.54	0.34	0.03	0.02
Detritus	32.07 ^a	28.23ª	8.25 ^b	4.18ª	3.02*	1.35 [*]
Macrophyte	15.33ª	29.44 ^b	1.91°	4.20 ^ª	0.33*	1.42*
Nematode	0.84 ^a	ND	0.06ª	ND	0.00*	0.00*
Fish egg	0.42 ^ª	0.40a	0.32ª	0.30 ^a	0.00*	0.00*
Digested muscle	0.42 ^ª	ND	0.11ª	ND	0.00*	0.00*

Table 5. Diet compositions of Oreochromis niloticus in Lake Hawassa on a rainy (n = 711) and dry (n = 248) months

^{a,b} Values of respective food items under the same category given different superscript letters are significantly different (α < 0.05).

Values of respective food items under the same category given different superscript letters are significantly different between rainy and dry months comparisons.-

ND, not detected.

Length-weight and length-length relationship of Oreochromis niloticus from Lake Hawassa

The length-weight relationship of *O. niloticus* in the present study showed a curvilinear relationship (Fig. 5A–5C), and the regression parameters generated from the power function of the length-weight relationship are statistically significant (ANOVA, p < 0.05). The slope of the regression line (b) for males, females, and mixed sexes of *O. niloticus* were 3.23, 3.25, and 3.24, respectively. The growth of both sexes followed a positive allometric growth pattern (b > 3). The *p*-value of the exponent (b) was significantly different from 3 for males, females, and both sexes (*t*-test, p < 0.001). The length-length relationship of *O. niloticus* showed a strong relationship ($r^2 = 0.99$), and is highly correlated (r = 0.99; Fig 5D).

Males: TW = 0.0077TL^{3,23}, $R^2 = 0.9841$, n = 504 Females: TW = 0.0073TL^{3,25}, $R^2 = 0.9849$, n = 450 Mixed: TW = 0.0075TL^{3,24}, $R^2 = 0.9844$, n = 954

Relative condition factor of *Oreochromis niloticus* from Lake Hawassa

The relative condition factors of *O. niloticus* from Lake Hawassa ranged from a minimum value of 0.52 to a maximum value of

1.58 for females with a mean \pm SE value of 1.01 \pm 0.0063, while that of the males ranged from 0.63 to 1.87 with a mean value of 1.08 \pm 0.0074. The largest relative condition factor was recorded in February and March for both sexes, while the smallest value was recorded in July for both sexes. The value of the relative condition factor indicates a significant variation between sampling months (ANOVA, p < 0.05). However, the interaction between months by sex shows an insignificant variation (ANOVA, p > 0.05), meaning the trend for the *O. niloticus* relative condition factor shows a similar fluctuation pattern between sexes (Fig. 6).

Discussion

Reproductive biology of *Oreochromis niloticus* from Lake Hawassa

The overall sex ratio (M:F) for *O. niloticus* in the present investigation was 1:0.93, test = 2.37, suggesting that, there was no significant difference from the theoretical 1:1 sex ratio. The current results are consistent with other research, which found no significant differences in the *O. niloticus* sex ratios such as from Lake Boyo as reported by Handago et al. (2024), and Lake Awassa as reported by Admassu (1994). However,



Fig. 5. The relationship between total length (TL) and total weight (TW) of (A) males, (B) females, both sexes (C), and TL and standard length (SL) (D) of *Oreochromis niloticus* from Lake Hawassa.



Fig. 6. The relative condition factors of *Oreochromis niloticus* from Lake Hawassa.

the current result showed contrary to certain other studies that found significant differences in the sex ratios of the same species such as: from Lake Tana as reported by Tadesse (1997), and from Lake Hayq as reported by Tessema et al. (2019). Sex ratio variations might be related to different reasons such as the method of fishing (Admassu, 1994), fishing pressure (Tekle-Giorgis et al., 2017), and sexual segregation (Dadebo et al., 2011).

In the current study, TL_{50} was 18.5 cm for males and 19.1

cm for females, respectively. The TL_{50} from the present study showed a larger value compared to other studies from the same species such as: from Lake Hayq where the TL_{50} value was 12.8 cm for males and 12.9 cm for females (Tessema et al., 2019), and 10.0 cm for males and 7.7 cm for females from Lake Boyo (Handago et al., 2024). However, compared to certain different lakes that were reported from the same species, the TL_{50} % maturity was smaller in Lake Hawassa. These included from the same study area Lake Hawassa 20.8 for males, and 20.3 for females O. niloticus (Muluve et al., 2016), and 20.5 cm for males and 19.6 cm for females from Lake Tana (Degsera et al., 2021). The variations of TL_{50} % maturity might be fishing pressure, mortality, and food availability (Admassu, 1994). For example, Muluye et al. (2016) reported 20.8 for females and 20.3 for males O. niloticus in the same study area, which indicates a higher value than the current research. However, studies from Lake Hawassa that focused entirely on dealing with problems related to fish stock assessments reveal a trend toward fishing that is becoming more than the safe level (Tekle-Giorgis, 2018). Harvesting of immature fish, rising fish mortality, and exploitation are negatively correlated with TL₅₀ (Tekle-Giorgis et al., 2017). Fish try to adapt to new environmental conditions and shift to sustain their recruits when the fishing pressure increases (Dadebo et al., 2011).

In the current investigation, O. niloticus breeds throughout

the year in Lake Hawassa, it was seen February through April is the first peak breeding season, and January is the second peak breeding season. The present finding agrees with different studies on the same species which were identified from different water bodies with the exceptions of some peak breeding months. For instance, O. niloticus breeds throughout the year in the same study area, with January to March and August to October representing the peak breeding seasons (Admassu, 1994). February to March, and July to September from Lake Boyo (Handago et al., 2024), between February and April in Lake Hayq (Tessema et al., 2019) were seen as the peak breeding months of O. niloticus. The variations of ripe fish proportions might be a result of temporal variation (Tadesse, 1997), and food availability (Admassu, 1996). O. niloticus prefers to reproduce where their offspring survive more especially during the rainy months (Admassu, 1996). Based on the present study, O. niloticus showed intensive breeding in both rainy and dry months. This might be the effect of limnological-related aspects. The diverse compositions of phytoplankton and zooplankton from Lake Hawassa help to continue their breeding at any time of the year when they get favorable situations (Admassu, 1996). Additionally, the dominant food resource of O. niloticus like blue-green algae specifically microsistis from Lake Hawassa didn't show a significant change in rainy and dry months (Kebede & Belay, 1994). The above-mentioned taxa of bluegreen algae are protein-rich (de la Fuente et al., 1977), and might create a favorable situation for O. niloticus breeding at any time of the year in Lake Hawassa.

With a mean value of 556 eggs per fish, the fecundity *O. niloticus* in Lake Hawassa ranged from 105 eggs per fish to the highest value of 1,541. *O. niloticus* from Lake Hawassa have good fecundity potentials in contrast to Lake Hayq, where 217 eggs per fish were reported (Tessema et al., 2019). However, *O. niloticus* from Lake Hawassa shows very poor fecundity potential than Lake Tana, where 730 eggs/fish on average (Tadesse, 1997). The individual size difference, ovary size, and quality of food resources might be the reason for variations in the fecundity potential of *O. niloticus*. For instance, the digestive efficiency of Lake Hawassa *O. niloticus* is higher than Lake Langano (Tadesse, 1999). Therefore, the quality of prey that they eat may limit their fecundity potential.

Food and feeding habits of *Oreochromis niloticus* from Lake Hawassa

The O. niloticus prey items found in Lake Hawassa are similar to

those seen in other lakes of the same species such as from Lake Tinishu Abaya (Anteneh et al., 2023), Lake Langano (Temesgen et al., 2022), Koka reservoir (Engdaw et al., 2013), and from the same study area (Tudorancea et al., 1988), and from the Gulf of Gorgora (Lake Tana) as reported by Engdaw (2023). Among the identified prey items of O. niloticus from Lake Hawassa, phytoplankton are the dominant (primary) prey item, which occurred at 95.72% of non-empty O. niloticus. This result agrees with some other Ethiopian Rift Valley Lakes which reported that phytoplankton were the primary prey for O. niloticus. For instance, the occurrences of phytoplankton from O. niloticus from Tinishu Abaya showed 94.1% (Anteneh et al., 2023), 100.0% from Lake Langano (Temesgen et al., 2022), 77% from Lake Tana (Engdaw, 2023), and 80.1% from Koka reservoir (Engdaw et al., 2013). Unlike the present finding, Handago et al. (2024) from Lake Boyo reported detritus was the most frequently occurring prey item, which occurred 99.2%.

Compared to the different taxa of phytoplankton, bluegreen algae are the most frequently occurring prey item from *O. niloticus* of Lake Hawassa based on the present finding (Table 3). Similar results have been reported from Lake Langano (Temesgen et al., 2022), from Tinishu Abaya (Anteneh et al., 2023), Gulf of Gorgora, (Engdaw, 2023), and from Koka Reservoir (Engdaw et al., 2013). Contradictory to the present finding, Tadesse (1999) from Lake Langano, reported that diatoms are the most frequently occurring prey. This variation might be related to the individual preference of fish, morphometry of Lake, biomass, and abundance of phytoplankton.

Based on volumetric contributions, phytoplankton contributed 88.2% of O. niloticus from Lake Hawassa and it is considered a primary prey item. The present finding agrees with other reports which indicate phytoplankton from non-empty O. niloticus contributed 39.5% from Tinishu Abaya (Anteneh et al., 2023), 47.5% from the Koka reservoir (Engdaw et al., 2013), 44.3% from the Gorgora Gulf of Lake Tana (Engdaw, 2023), and 64.3% from Lake Langano (Temesgen et al., 2022). Unlike the present finding, the volumetric contributions of macrophytes from Lake Boyo (73.6%) were the primary prey items for O. niloticus (Handago et al., 2024). Detritus is the second most occurred prey item of O. nilotics in both frequency of occurrence and volumetric contribution accounting for 31.1% and 7.2% from Lake Hawassa. This finding agrees with other Ethiopian Rift Valley Lakes of O. niloticus prey items, such as O. niloticus from Lake Tinishu Abaya detritus where the second

important prey item occurred at 74.3% (Anteneh et al., 2023). Contradictory from the present result, macrophyte was the second contributing prey (25.8%) in volumetric contribution from Lake Tinishu Abaya (Anteneh et al., 2023).

Cladocera is a major prey item, compared to other texa of zooplankton which occurred at 7.9%. Unlike the present finding, rotifers were a major prey item compared to other zooplanktons texa from other Rift Valley Lakes such as O. niloticus from Lake Langano (Temesgen et al., 2022), and Tinishu Abaya (Anteneh et al., 2023). This might be the effect of pollution where the biomass of zooplankton including Rotifera from Lake Hawassa shows a decreasing trend as a result of anthropogenic effects (Beyene et al., 2022). The occurrence and contributions of nematode from Lake Hawassa are too low compared to Lake Tana which reported 28.1%, and 0.1% respectively (Engdaw, 2023), and from Lake Langano which indicates 7.8% and 0.3% respectively (Temesgen et al., 2022). These variations might be related to the protein source of prey items (Tadesse, 1998). In the case of Lake Hawassa, there are different optional planktons contributed as protein sources. Additionally, some algae contributed as a prey item for O. niloticus in the present study such as blue-green algae specifically microsists are the most frequent prey item compared to different blue-green algae. Microsists are proteinrich (de la Fuente et al., 1977), and O. niloticus concentrates on protein-rich prey (Tadesse, 1998).

Based on the index of preponderance methods of comparison, the major prey item for *O. niloticus* from Lake Hawassa showed 96.6% phytoplankton (Table 3). A similar finding is reported from other inland water bodies of Ethiopia. For example, Temesgen et al. (2022) from Lake Langano reported 74.5%, and 52.3% were reported from Lake Tana (Engdaw, 2023). Contradictory to the present finding, detritus denotes 79.1% from Lake Boyo (Handago et al., 2024).

Unlike the present finding, it was reported that aquatic insects are considered important prey items for *O. niloticus*. For instance, from Lake Tana (Engdaw, 2023), Lake Langano (Temesgen et al., 2022), and the same study area Lake Hawassa, zoobenthos also has a contribution as a prey for *O. niloticus* (Tudorancea et al., 1988). These variations might be the effect of interspecific competitions. Aquatic insects from Lake Hawassa have a role as a diet of larger fish species of Lake Hawassa like *C. gariepinus* (Dadebo, 2000). The present finding indicates there was a slight ontogenetic dietary shift of *O. niloticus* from Lake Hawassa (Fig. 4) but not significance shift (C > 0.6). In all

size classes, the contributions of phytoplankton were identified from the stomach of *O. niloticus* (Fig. 4). The contributions of animal-origin sources are inversely related to the size class of *O. niloticus* from Lake Hawassa. A similar finding was reported from Koka Reservoir (Engdaw et al., 2013). This indicates that in lower-size classes, fish require a high protein source for survival and overall growth activities (Temesgen et al., 2022). On the other hand, *O. niloticus* in the lower size class the size of their stomach is small, and have not much space to load and digest larger food items like plant origins (Engdaw et al., 2013).

At all, except for the volumetric contributions of detritus and occurrences of macrophyte, there was no statistical significance diet composition of *O. niloticus* in Lake Hawassa from the present study on rainy and dry seasons comparisons (Table 5). The present finding contradicts other studies that were carried out at Lake Langano (Temesgen et al., 2022). The above-mentioned researchers identified a seasonal variation in phytoplankton occurrences from the stomach contents of *O. niloticus*. The present non-significance phytoplankton variations might be the effect of dietary overlap. The present study indicates a strong overlap index (Table 4) and the preference of prey might be limited.

Length-weight and length-length relationship of Oreochromis niloticus

Based on the current study, O. niloticus has a curvilinear length-weight relationship and is highly significant ($r^2 > 0.98$, p < 0.05). The finding agrees with other studies on the same species such as from Lake Boyo (Handago et al., 2024), and Lake Tana (Engdaw, 2023). The finding also confirms some former works which were identified as allometric growth such as in Lake Boyo with the value of regression coefficient 3.18 for females, and 3.12 for combined sexes exhibited slight positive allometric growth pattern as reported by (Handago et al., 2024). The present results are in disagree with some findings that were identified as isometric, and negative allometric growth, such as those from Lake Hayq, where b = 2.50 for females and 2.33 for males, respectively (Tessema et al., 2019), and b = 2.8 from the Gulf of Gorgora in Lake Tana (Engdaw, 2023). These variations of slope (b) of O. niloticus depend on several external and internal factors. During the present study, the dry months were very short. This might affect the total biomass production in the lake which might create good conditions for growth. There is a contradictory idea of the primary productions of Lake Hawassa. for instance, primary production increases in wet months compared to rainy months (Gebre-Mariam & Taylor, 1989). In a contradictory way, the nutrient mixing and primary production from Lake Hawassa increase during the dry month (Tilahun & Ahlgren, 2010). This situation might be helping the growth of *O. niloticus* to get good food resources in both dry and rainy months.

Relative condition factors of Oreochromis niloticus

The mean relative condition factors of *O. niloticus* for the present study were found to be 1.08 ± 0.0074 for males and 1.01 ± 0.0063 for females. The mean relative condition factors for both sexes showed K_n > 1, indicating that the health status of *O. niloticus* from Lake Hawassa is currently safe (Jones et al., 1999). These findings align with comparable values reported for *O. niloticus* populations in Lake Victoria (Njiru et al., 2006). The slight differences in relative condition factors between males and females may reflect physiological demands related to reproduction and environmental influences during the sampling months (Jones et al, 1999).

Conclusion

The present study shows that O. niloticus male-to-female sex ratio (1:0.93) is almost equal to the theoretical 1:1 ratio. The breeding seasons of O. niloticus from Lake Hawassa were bimodal, intensively breeding from February to April, and by January. O. niloticus under 19.1 cm total length should be protected to save the non-breeding fish from unwise fishing. It is also recommended to close at the peak breeding season for continuing their recruits. Based on the stomach contents of O. niloticus from Lake Hawassa, it is seen that both plant and animal origin are ingested. Strong dietary overlaps are observed in the present finding and are predominantly focused on phytoplankton. Insignificant dietary shift and phytoplankton dominance in the diet of O. niloticus suggest that a parallel study on phytoplankton biomass and primary production, as a proxy of yield and management of Tilapia stock in Lake Hawassa, should be considered for future studies.

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was funded and supported by the Ethiopian Fed-

eral Ministry of Education (517 USD) and Dilla University (431 USD) for data collection. All materials and laboratory access were sponsored by the Department of Aquatic Sciences, Fisheries, and Aquaculture at Hawassa University, Ethiopia.

Acknowledgements

The authors acknowledge the department of Aquatic Sciences, Fisheries, and Aquaculture, College of Natural and Computational Sciences, Hawassa University for granting laboratory and field materials. We also thank the Center for Aquaculture Research and Education (CARE) of Hawassa University for material support during the field work.

Availability of data and materials

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Ethics approval and consent to participate

This study does not involve the use of human subjects (human-related experimentation) but involves field sampling of fish from non-protected sites. However, ethical clearance to carry out field observation and laboratory work was ethically granted by Hawassa University College of Natural and Computational Sciences under reference number CNCS-REC012/23.

ORCID

Teshome Belay Elias Dadebo	https://orcid.org/0000-0001-7072-0060
Girma Tilahun	https://orcid.org/0000-0000-0000-0000
Degsera Aemro	https://orcid.org/0009-0000-9781-6011 https://orcid.org/0000-0003-0073-4111

References

Admassu D. Maturity, fecundity, brood-size and sex ratio of tilapia (*Oreochromis niloticus* L.) in Lake Awassa. SINET Ethiop J Sci. 1994;17:53-69.

Admassu D. The breeding season of tilapia, *Oreochromis niloticus* L. in Lake Awassa (Ethiopian rift valley). Hydrobiologia. 1996 ;337:77-83.

Anteneh YE, Mamo SW, Tilahun MA. Fisheries in Lake Tinishu Abaya (Ethiopia) could be managed using dietary nature of Nile tilapia (*Oreochromis niloticus* L. 1757). Fish Aquat Sci. 2023;26:491-9.

- Beyene G, Kifle D, Fetahi T. Spatial distribution of zooplankton in relation to some selected physicochemical water quality parameters of Lake Hawassa, Ethiopia. Afr J Aquat Sci. 2022;47:163-72.
- Bowen, SH. Quantitative description of the diet. 2nd ed. Bethesda, MD: American Fisheries Society; 1996.
- Dadebo E. Reproductive biology and feeding habits of the catfish *Clarias gariepinus* (Burchell) (Pisces: Clariidae) in Lake Awassa, Ethiopia. SINET Ethiop J Sci. 2000;23:231-46.
- Dadebo E, Gebre-Mariam Z, Mengistou S. Breeding season, maturation, fecundity and condition factor of the African catfish *Clarias gariepinus* Burchell 1822 (Pisces: Clariidae) in Lake Chamo, Ethiopia. Ethiop J Biol Sci. 2011;10:1-17.
- Dadebo E, Kebtineh N, Sorsa S, Balkew K. Food and feeding habits of the red-belly tilapia (*Tilapia zillii* Gervais, 1848) (Pisces: Cichlidae) in Lake Ziway, Ethiopia. Agric For Fish. 2014;3:17-23.
- Dadebo E, Tadele B, Balkew K. The impact of gillnet selectivity on immature Nile tilapia (*Oreochromis niloticus* L.) (Pisces: Cichlidae) in LakeHawassa, Ethiopia. In: Proceedings of the EFASA Fifth Annual Conference; 2023; Hawassa, Ethiopia. p. 79-96.
- Degsera A, Minwyelet M, Yosef TG. 2021. Stock assessment of Nile tilapia *Oreochromis niloticus* (Linnaeus 1758) in Lake Tana, Ethiopia. Afr J Aquat Sci. 2021;46:499-507.
- de la Fuente G, Flores A, Molina MR, Almengor L, Bressani R. Some nutritional characteristics of a naturally occurring alga (*Microcystis* sp.) in a Guatemalan Lake. Appl Environ Microbiol. 1977;33:6-9.
- Demsie AF, Yimer GT. Occurrence of microplastics in commercial fish species from the Ethiopian rift valley's Lake Hawassa, Ethiopia. Environ Sci Pollut Res. 2024;31:48641-9.
- Engdaw F. Morphometric relations and diet compositions of Nile tilapia *Oreochromis niloticus* (Linn. 1758) in Lake Tana Gorgora gulf, Ethiopia. Fish Aquat Sci. 2023;26:169-80.
- Engdaw F, Dadebo E, Nagappan R. Morphometric relationships and feeding habits of Nile tilapia *Oreochromis niloticus* (L.) (Pisces: Cichlidae) from Lake Koka, Ethiopia. Int J Fish Aquat Sci. 2013;2:65-71.

- Getachew T, Fernando CH. The food habits of an herbivorous fish (*Oreochromis niloticus* Linn.) in Lake Awasa, Ethiopia. Hydrobiologia. 1989;174:195-200.
- Gebre-Mariam Z, Taylor WD. Seasonality and spatial variation in abundance, biomass and activity of heterotrophic bacterioplankton in relation to some biotic and abiotic variables in an Ethiopian rift-valley lake (Awassa). Freshw Biol. 1989;22:355-68.
- Handago T, Dadebo E, Tilahun G. Reproductive biology and feeding habits of Nile tilapia *Oreochromis niloticus* (Linnaeus, 1758) (Pisces: Cichildae) in Lake Boyo, Ethiopia. Ethiop J Sci Technol. 2024;17:81-97.
- Holden MJ, Raitt DFS. Manual of fisheries science: part 2 methods of resource investigation and their application.
 FAO Fisheries Technical Paper No. 115. Rome: Food and Agriculture Organization (FAO); 1974.
- Hyslop EJ. Stomach contents analysis—a review of methods and their application. J Fish Biol. 1980;17:411-29.
- Jones RE, Petrell RJ, Pauly D. Using modified length-weight relationships to assess the condition of fish. Aquacult Eng. 1999;20:261-76.
- Kebede E, Belay A. Species composition and phytoplankton biomass in a tropical African lake (Lake Awassa, Ethiopia). Hydrobiologia. 1994;288:13-32.
- King M. Fisheries biology, assessment and management. Oxford: John Wiley & Sons; 2013.
- Muluye T, Tekle-Giorgis Y, Tilahun G. The extent of immature fish harvesting by the commercial fishery in Lake Hawassa, Ethiopia. Momona Ethiop J Sci. 2016;8:37-49.
- Natarajan AV, Jhingran AG. Index of preponderance— a method of grading the food elements in the stomach analysis of Fishes. Indian J Fish. 1961;8:54-9.
- Njiru M, Ojuok JE, Okeyo-Owuor JB, Muchiri M, Ntiba MJ, Cowx IG. Some biological aspects and life history strategies of Nile tilapia *Oreochromis niloticus* (L.) in Lake Victoria, Kenya. Afr J Ecol. 2006;44:30-7.
- Pauly D. Fish population dynamics in tropical waters: a manual for use with programmable calculators. Penang: WorldFish; 1984.
- Reyntjens D, Wudneh T. Fisheries management a review of the current status and research needs in Ethiopia. SINET Ethiop J Sci. 1998;21:231-66.
- Schoener TW. Nonsynchronous spatial overlap of lizards in patchy habitats. Ecology. 1970;51:408-18.
- Tadesse Z. Breeding season, fecundity, length-weight rela-

tionship and condition factor of *Oreochromis niloticus* L. (Pisces: Cichlidae) in Lake Tana, Ethiopia. SINET Ethiop J Sci. 1997;20:31-47.

- Tadesse Z. Food and feeding ecology of tilapia, *Oreochromis niloticus* L. and effects of diet on the lipid quality of fish in some lakes in Ethiopia. Addis Ababa: Addis Ababa University; 1998.
- Tadesse Z. The nutritional status and digestibility of *Oreochromis niloticus* L. diet in Lake Langeno, Ethiopia. Hydrobiologia. 1999;416:97-106.
- Tekle-Giorgis Y. Prediction of bio-economically sustainable yield and optimum fishing effort for the Nile tilapia (*Oreochromis niloticus* L.) fishery of Lake Hawassa, Ethiopia. Ethiop J Sci Technol. 2018;11:57-78.
- Tekle-Giorgis Y, Berihun A, Dadebo E. Assessment of sustainable yield and optimum fishing effort for the tilapia (*Oreochromis niloticus* L. 1758) stock of Lake Hawassa, Ethiopia. Momona Ethiop J Sci. 2017;9:1-21.
- Tekle-Giorgis Y, Wagaw S, Dadebo E. The food and feeding habits of the African catfish, *Clarias gariepinus* (Burchell, 1822) (Pisces: Clariidae) in Lake Hawassa and Shallo swamp, Ethiopia. Ethiop J Biol Sci. 2016;15:1-18.
- Temesgen M, Getahun A, Lemma B, Janssens GPJ. Food and feeding biology of Nile tilapia (*Oreochromis niloticus*) in Lake Langeno, Ethiopia. Sustainability. 2022;14:974.
- Tesfaye G, Wolff M. The state of inland fisheries in Ethiopia: a synopsis with updated estimates of potential yield. Ecohydrol Hydrobiol. 2014;14:200-19.
- Tessema A, Getahun A, Mengistou S, Fetahi T, Dejen E. Length-weight relationship, condition factor and some reproductive aspects of Nile tilapia (*Oreochromis niloticus*) in Lake Hayq, Ethiopia. Int J Fish Aquat Stud. 2019;7:555-61.
- Tilahun G. Waste management in Hawassa city: anthropogenic threat on Lake Hawassa. Sustain Syst. 2023;5.
- Tilahun G, Ahlgren G. Seasonal variations in phytoplankton biomass and primary production in the Ethiopian Rift Valley lakes Ziway, Awassa and Chamo – the basis for fish production. Limnologica. 2010;40:330-42.
- Tudorancea C, Fernando CH, Paggi JC. Food and feeding ecology of *Oreochromis niloticus* (Linnaeus, 1758) juveniles in Lake Awassa (Etiopia). Arch Hydrobiol Suppl Monogr Beitr. 1988;79:267-89.
- Zar J. Data transformations. 4th ed. Hoboken, NJ: Upper Saddle River; 1999.