



# Mapping and assessing the coral reef health in Kadatua, Tiwutonkidi and Siompu Islands, South East Sulawesi, Indonesia

Daniel Deonisius Pelasula<sup>1, 2, 3, \*</sup>, Sam Wouthuyzen<sup>2, 3</sup>, Mufti Petala Patria<sup>1</sup>, Simon Izaak Patty<sup>2</sup>, Petrus Christianus Makatipu<sup>2</sup>, Baru Sadarun<sup>4</sup>, Wiesye Violent Pelupessy<sup>5, 6</sup>, Suiyuan Wang<sup>6, 7</sup>, Johanis Dominggus Lekallete<sup>3, 8</sup>

<sup>1</sup> Faculty of Mathematics and Natural Sciences, University of Indonesia, Depok 16424, Indonesia

<sup>2</sup> Research Centre for Oceanography, National Research and Innovation Agency, Jakarta 14430, Indonesia

<sup>3</sup> Centre for Collaborative Research on Aquatic Ecosystems in Eastern Indonesia, Ambon 97233, Indonesia

<sup>4</sup> Faculty of Fisheries and Marine Sciences, Halu Oleo University, Kendari 93232, Indonesia

<sup>5</sup> Faculty of Engineering, Pattimura University, Ambon 97233, Indonesia

<sup>6</sup> Department of Urban and Regional Planning, University at Buffalo, New York, NY 14260, USA

<sup>7</sup> Department of Geography, University at Buffalo, New York, NY 14261, USA

<sup>8</sup> Research Center for Climate and Atmosphere, National Research and Innovation Agency, Bandung 40135, Indonesia

## Abstract

The quality and quantity of coral reefs contain essential ecological functions to provide goods and environmental services for the livelihood of local communities. However, coral reefs continue to suffer in most of Indonesia's remote reefs. This study aims to map and monitor the health coral reef in five sites of south Buton Regency, the Kadatua (BTNC-13 and -14), Tiwutonkidi (BTNC-C), and Siompu Islands (BTNC-D and -17) using the effective and efficient remotely sensed Landsat-8 OLI (operational land imager) satellite and intensive field survey data using Line intercept transect (LIT) method, then use the result to assess the coral health index (CHI) of the study sites. The results of the mapping using Landsat-8 OLI data classified five groups of benthic habitats: algae, corals, seaweed, rubble, and sand, after following the standard image processing procedure and utilizing sun-glint, water column correction and maximum likelihood classification. Meanwhile, the intensive LIT survey resulted in information on the percent live coral cover that was in the range of moderate conditions (25 –≤ 50%), but the site of BTNC-17 was in bad condition (< 25%). The resilient condition of coral reefs in all sites was good, which is indicated by the low cover of fleshy seaweed but the low biomass of herbivorous and carnivorous reef fish. Thus, the CHI, in general, was in moderate condition (CHI 5 and 6). However, a comparison with CHI data from 2016 showed that the coral condition in study sites tends to decline. Several efforts to improve the CHI and to ensure the long-term health and resilience of coral reef ecosystems are suggested. The findings of this study can be used as a benchmark and indicator to monitor the progress or the decline of the coral reef environment in the future at five study sites

Received: Oct 8, 2024 Revised: Dec 24, 2024 Accepted: Jan 20, 2025

\*Corresponding author: Daniel Deonisius Pelasula

Faculty of Mathematics and Natural Sciences, University of Indonesia, Depok 16424, Indonesia

Tel: +62-811-4703-246, E-mail: pelasuladaniel@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2025 The Korean Society of Fisheries and Aquatic Science

in Kadatua, Tiwutunkidi and Siompu Islands.

**Keywords:** Landsat-8 OLI (operational land imager), Line intercept transect (LIT), Reef fish abundance, Coral health index (CHI), South of Buton Regency

## Introduction

Coral reefs, although they cover less than 1% of the Earth's surface and 2% of the seabed, support at least 25% of marine biodiversity (Spalding et al., 2001). They provide essential benefits to humanity, including maintaining the health of coastal ecosystems by facilitating important activities such as grazing, breeding, and spawning (Moberg & Folke, 1999), coastal protection (Ferrario et al., 2014), support for sustainable fisheries (Darling & D'agata, 2017), and contributions to marine tourism (Spalding et al., 2017). Additionally, coral reefs offer medicinal compounds (Cooper et al., 2014) and serve as natural laboratories for education and conservation sites (ICRI, 2020). In Indonesia, coral reefs hold significant economic value, supporting millions of communities across numerous small islands and coastal regions.

However, coral reefs are among the most vulnerable ecosystems. Nearly 20% of the world's coral reefs are in severe decline, with estimates suggesting that 30% could experience massive damage by 2032, and many may be lost by 2050 (Hughes et al., 2017). Meanwhile, the coral reef in Indonesia ranks first globally with 18% of the world's coral reefs, covering approximately 51,000 km<sup>2</sup>, primarily located in the coral triangle, the heart of the coral reefs of the world, which boasts 83 genera and 569 species of coral (Hoeksema, 2007). However, many reefs face degradation due to declines in environmental quality (Burke et al., 2012; Pandolfi et al., 2003). In Indonesia, a study by Suharsono (unpublished; see Slide 1 coraltriangleinitiative.org) revealed that out of 1,064 observation sites, 35.2% were in poor condition (live coral cover [LCC] < 25%), 35.1% were in fair condition (LCC 25 –< 50%), 23.4% were in good condition (LCC 50 –< 75%), and only 6.4% were in excellent condition (LCC > 75%). Wulandari et al. (2022) also found poor coral health in the Selayar Biosphere Reserve at Taka Bonerate Islands.

The deterioration of coral reefs, on a local scale, is caused by various anthropogenic stressors. These include overfishing and destructive fishing practices (Jackson et al., 2001), the impact of land-based pollution such as nutrient runoff, sedimen-

tation, and plastic pollution (Fabricius, 2005; Syakti et al., 2019), biological factors like predator population blooms of *Acanthaster planci* and *Drupella* spp snails (Brodie & McElroy, 2015; Kiel & Klemens, 2006), and coral diseases also play a significant role (Sutherland et al., 2004). On a global scale, climate change has led to ocean acidification and rising sea temperatures globally, causing coral bleaching (Hoegh-Guldberg et al., 2007; Hughes et al., 2017; Wouthuyzen et al., 2018). Those factors collectively contribute to the decline of coral reef ecosystems.

Coral reefs can recover from damage, but recovery times vary. Some species may heal in just 4 to 5 years, like those in the Marine Recreation Park of Pieh Island of west Sumatra (Wouthuyzen et al., 2019), while others may take up to 16 years (Fox et al., 2019). Therefore, mapping, monitoring, and evaluating coral reef conditions using remotely sensed satellite data is inevitable.

Various Earth observation satellites with high-resolution multispectral sensors, but those are must be purchased such as IKONOS (4 m), GeoEye-1 (1.65 m), Quickbird (1.61 m), WorldView-3 (1.24 m), and Pléiades Neo (30 cm), along with free medium-resolution options like Landsat-8/9 (30 m) and Sentinel-2 (10 m), are available for effective remote sensing. For instance, Nurdin et al. (2015) mapped coral reefs and their damages caused by destructive fishing practices using multi-temporal and multi-sensor Landsat satellite imagery in the Spermonde Islands, South Sulawesi, Indonesia, by applying image processing procedures similar to this study. Meanwhile, Aulia et al. (2020) mapped the changes of bottom benthic profile of the shallow-water coral reefs in Karya, Semak Daun Islands, and Balik Layar Reef in Seribu Islands, Indonesia, in 2016 and 2018 based on Landsat 8 (OLI) satellite imagery, which also used the same procedures as this study.

Coral reef mapping in Indonesia has utilized medium-resolution satellite data. Notable studies include Adji (2014), who mapped reefs in the Wakatobi Islands using Landsat-7. Haya & Fujii (2017) focused on the Spermonde Islands with multi-temporal Landsat data. Kartikasari et al. (2021) mapped reefs in Lovina, northern Bali, using Sentinel-2 data. However, detailed

information on coral reefs in Kadatua, Liwutonkidi, and Siompu Islands in South Buton Regency, which are crucial for local livelihoods dependent on fisheries and tourism, remains lacking. The Indonesian Ministry of Marine Affairs reported 471 cases of coral damage from harmful fishing practices in South and Southeast Sulawesi from 2013 to 2019, with Kadatua, Liwutonkidi and, and Siompu Island among the affected. This study aims to map the coral reef and develop the coral reef health at five sites in these islands, aiding local coral reef management and contributing to the World Coral Reef Status Database 2025 by the Global Coral Reef Monitoring Network (GCRMN).

## Material And Methods

### Study sites

This study was conducted on Kadatua, Liwutonkidi, and Siompu Islands in South Buton Regency, Southeast Sulawesi, Indonesia, located between latitudes  $-5.5^{\circ}$  to  $-5.7^{\circ}$  South and longitudes  $122.385$  to  $122.585^{\circ}$  East. Field surveys were carried out at five locations: two on Kadatua Island (BTNC 13, BTNC 14), one on Liwutonkidi Island (BTNC C), and two on Siompu Island (BTNC 17, BTNC D), as shown in Fig. 1.

The study utilized Landsat-8 OLI satellite imagery (Path 122, Row 064) acquired in January 13, 2021. Data was obtained from the USGS website at <https://earthexplorer.usgs.gov/>. Data collection involved SCUBA diving gear, a handheld Garmin

GPS, an underwater camera, and a diving slate with waterproof paper and pencils. Data processing was done using QGIS, a free and open-source GIS software.


### Satellite data analysis procedures

Satellite data analysis procedures involve assessing coral reef habitat coverage and evaluating the coral reef resilience and health. The process includes two stages: pre-image and image processing. During pre-image processing, steps such as atmospheric correction, image cropping, de-glint (sun glint correction), and masking clouds and land are performed to enhance image quality (Fig. 2).


The radiometric data from Landsat-8 OLI images were corrected using the supplied metadata by converting Digital Number (DN) values into radiance or reflectance. Atmospheric influences were adjusted using the Dark Object Subtraction (DOS) method (Song et al., 2001), and the images were cropped to match the specific study sites.

To improve Landsat data quality, a sun glint correction, also known as de-glint, was applied by selecting representative pixels for different sun glint visibility levels in bands 2, 3, 4, and 5 (blue, green, red, and near-infrared [NIR]). This de-glint process follows Hedley et al. (2005) as described in Equation (1).

$$B_{\text{corrected}} = B_i - b \times (NIR - NIR_{\min}), \quad B_i = a + b \times NIR \quad (1)$$



**Fig. 1.** The map shows the study site with five observation locations on the Kadatua (BTNC-13 and -14), Liwutonkidi, (BTNC-C) and Siompu Islands (BTNC-17 and -D).



**Fig. 2. Flow chart of analysis of the Landsat 8 (OLI) data to generate the benthic habitat map.**

where:  $B_{corrected}$  = de-glnt corrected visible band blue, green, or red of Band-2, -3, and -4, respectively.

$B_i$  = Landsat 8 OLI's Visible Band 2, 3 and 4;

$a$  and  $b$  are the intercept and slope of the regression line, respectively.

After the de-glnt process, we mask the study object to isolate it from unrelated features like clouds and land. This is followed by water column correction and supervised classification using maximum likelihood (MLH) classification. The depth invariant index (DII), based on Lyzenga (1981), employs a green-red (B2-B3) band combination from the Landsat 8 OLI image to capture underwater details accurately. This correction ensures precise information about underwater objects follows the methods outlined by Green et al. (2000) and Lyzenga (1981) using the equation;

$$DII = \ln(L_i) - \left( \frac{K_i}{K_j} \right) \times \ln(L_j) \quad (2)$$

(3)

$$\left( \frac{K_i}{K_j} \right) = a + \sqrt{(a^2 + 1)} \quad (3)$$

$$a = (\text{var}_i - \text{var}_j) / 2 \times \text{covar}_{ij} \quad (4)$$

where DII is the depth Invariant Index,

$L_i$  is the digital value in band  $i$ ,  $L_j$  is the digital value in band  $j$ ,

$K_i/K_j$ : is the ratio of the attenuation coefficient in the band pairs  $i$  and  $j$ ,

$\text{Var}_i$ : is variant band  $i$ ;  $\text{Var}_j$ : is variant band  $j$ ;  $\text{Covar}_{ij}$ : is covariant band  $ij$ .

The attenuation coefficient is calculated by defining a region of interest (ROI) in a homogeneous area, with measurements taken at various depths. The ROI is identified through visual interpretation and underwater photography. After correcting for the water column, image classification is performed using MLH classification, which categorizes pixels based on their probability of belonging to specific classes. A pixel is not classified if its value falls below a defined threshold. Ground truthing at each data site was conducted at 62 sites using a Garmin GPS device, following SNI 7716-2011 guidelines for mapping demersal habitats in shallow waters (LIPI, 2014). Accuracy testing was done with a confusion matrix comparing classification results to field data (Congalton & Green, 2019).

### Coral reef data collection

To collect benthic habitat data on coral reefs, underwater photos were taken using the Photo Transect method (Giyanto & Soedarma, 2010). Observers equipped with SQUBA diving gear, underwater cameras, and bright yellow rectangular frames (58 × 44 cm) took pictures positioned 60 cm above the substrate along a 50-meter transect line (Fig. 3A). A total of 50 photos, 25 from the left and 25 from the right side along the transect line were taken.

The analysis of fifty underwater photos (Fig. 3B) utilized Coral Point Count with Excel extensions (CPCe®), developed by Nova Southeastern University (Kohler & Gill, 2006). We applied a benthic component coverage technique by randomly selecting 25 points per photo and categorizing them using the CPCe® database. After exporting the results to Microsoft Excel, we calculated the percentage coverage of benthic components to assess the CHI.

### Coral reef fish abundance and biomass

Coral reef fish observations were conducted using the underwater visual census (UVC) method from the ASEAN-Australia Project, which allows for rapid estimates of fish abundance, biomass, and distribution (Rolim et al., 2022). This assessment helps evaluate coral reef health. The UVC involved a 70-meter transect line with a 2.5-meter observation area on each side, thus resulting in a total area of 350 m<sup>2</sup> per transect, as shown in Fig. 4. The standing stock ( $S$ ) of each reef fish group was calculated using the following equation:






Fig. 3. Underwater photo transect (A) and photo transect analysis according to CPCe<sup>®</sup>, tool (B).




Fig. 4. Illustration of reef fish stock assessment using underwater visual census (UVC).

$S$  (Fish number) =  $A$  (Coral reef area: Ha)  $\times$   $D$  (Fish Density, fish/ha) (5)

Coral health index (CHI) is assessed based on the biomass density of reef fish, categorized into two groups: (1) Carnivorous fish (Families: Haemulidae, Lethrinidae, Lutjanidae, Serranidae), and (2) Herbivorous fish (Families: Acanthuridae, Scaridae, Siganidae). Thus, CHI was evaluated by converting the fish counts (numbers) of those groups into biomass density (B: kg/ha) using the equation:

$B$  (Total Fish Biomass, kg/ha) =  $\sum [S$  (Number of a certain fish species)  $\times W$  / Reef areas (ha)] (6)

where,  $S$  = Fish Number of a certain species;

$W = a L^b$ ,  $W$ : is Weight of a certain fish species (gram or kg),  $L = TL$ : is the total length of the fish (cm),  $TL = (L_{min} + L_{max})/2$  or  $TL = 0.65 \times L_{max}$ , whereas  $L_{min}$  and  $L_{max}$  are the minimum and maximum fish TL that can be obtained from Froese & Pauly (2018);

$a$  and  $b$ : is the fish growth coefficient that can be cited from Kulbicki et al. (2005).

### Coral health index

The CHI comprises several fundamental components, including the percentage of LCC, fleshy seaweed (FS) cover as indicator of recovery potential or resilience levels, and coral reef fish biomass (RFB) (Giyanto et al., 2017). The specific criteria for evaluating the coral reef's health of these components are detailed in Table 1. Meanwhile, the CHI is assessed using those

Table 1. Criteria for assessing the coral healthy index (CHI)

No	Category	Coral cover criteria	Recovery potential or resilience levels	Fish Biomass criteria
1.	Low	< 19%	Fleshy seaweed cover > 3%, coral fragment cover > 60%, and live coral cover < 5%	< 970 kg/ha
2.	Moderate	19–35%	—	970–1,940 kg/ha
3.	High	>35%	Fleshy seaweed cover <3%, coral fragment cover < 60%, and live coral cover >5%	>1,940 kg/ha

Data from Giyanto et al., (2017).

**Table 2. CHI based on LCC, Fleshy seaweeds /recovering potential, and coral fish components**

No	Benthic components		Coral fish components	
	Live coral cover (LCC)	Fleshy seaweed cover or Resilience potential	Reef fish biomass (RFB)	Coral health index (CHI)
1	High	High	High	10
2	Moderate	High	High	9
3	High	High	Moderate	8
4	High	Low	High	8
5	Moderate	High	Moderate	7
6	Low	High	High	7
7	High	High	Low	6
8	High	Low	Moderate	6
9	Moderate	Low	High	6
10	Moderate	High	Low	5
11	Low	High	Moderate	5
12	Low	Low	High	5
13	High	Low	Low	4
14	Moderate	Low	Moderate	4
15	Low	High	Low	3
16	Low	Low	Moderate	3
17	Moderate	Low	Low	2
18	Low	Low	Low	1

Data from Giyanto et al., (2017).

components, resulting in 18 combinations of CHI values that range from 10 to 1 as detailed in Table 2. An index level of 10 in Table 2 indicates the healthiest coral reefs, characterized by high coral cover and resilience, which supports diverse coral fish and higher fish biomass. Conversely, an index value of 1 signifies poor coral cover and low resilience, hindering coral recovery and growth and correlating with very low fish biomass due to the degraded ecosystem that cannot sustain coral fish.

## Results

### Result of the processing of Landsat-8 Oli image

The Landsat-8 OLI satellite image data was pre-processed to enhance clarity and minimize atmospheric interference. This interference can cause the minimum DN value for the darkest objects, such as cloud shadows and deep sea, to be zero (0). Table 3 shows a minimum DN value of 0 after atmospheric correction using the DOS method, with maximum values in most bands exceeding 100, which often occur due to a few pixels affected by thick clouds. This correction is essential for reducing atmospheric interference. Landsat-8 has 11 bands, with 30 m resolution for multispectral bands (1–7) and 15 m for the pan-


**Table 3. Statistical results of atmospheric corrections**

Bands	Min	Max	Mean	StdDev
Band 2 (Blue)	0.0000	144.1036	9.7764	11.7012
Band 3 (Green)	0.0000	144.1036	7.0298	11.2047
Band 4 (Red)	0.0000	144.1036	5.6439	11.4764
Band 5 (NIR)	0.0000	144.1036	7.5808	15.6917

NIR, near-infrared.

chromatic band. In this study, we analyzed four visible bands: the Blue (Band 2), Green (Band 3), Red (Band 4), and near-infrared/NIR (Band 5) (Table 3).

Coral reefs, seagrass, and algae habitats are often below the sea surface, making accurate identification from satellite imagery challenging due to atmospheric interference and water column effects (Chen et al., 2021). Sunlight reflection and ripples can blur these features, necessitating the correction of sun glint (Fell, 2022). In this study, sun glint was corrected using a linear regression equation that relates the Blue, Green, and Red bands to the NIR band (Hedley et al., 2005). Fig. 5 illustrates the relationship between the Blue, Green, and Red bands to the NIR bands. The minimum NIR value (NIRmin) extracted from the NIR band (Band 5) was 1.4918. The regression equation,



**Fig. 5. Relationship between Band 2, Band 3, and B 4 versus B 5 for assessing the sun-glint.**

such as that given by Equation (1) ( $Bi_{corrected} = Bi - bi \times [NIR - NIR_{min}]$ ), is shown in Table 4. This corrected image was then used for water column attenuation correction.

Water column correction is generally used to eliminate the fading effect on an object's signal as it travels through the water column caused by the attenuation of electromagnetic waves (Lyzenga, 1981). This attenuation leads to varying pixel values for the same object at different depths, with reflectance values decreasing as depth increases (Green et al., 2000). Thus, the correction for each band is determined using a linear regression equation based on combined visible light bands, where the slope represents the attenuation coefficient ratio (Lyzenga, 1981). Sand is chosen for sampling because it is easily identifiable at various depths; the plots of band pairs of Band 2 versus Band 3,


Band 2 versus Band 4, and Band 3 versus Band 4 are shown in Fig. 6.

The determination coefficients ( $R^2$ ) of Fig. 6A–6C are considered sufficiently high to warrant further analysis. The  $R^2$  values for the Band pairs of  $\ln(B2)$  vs  $\ln(B3) = 0.9608$ ;  $\ln(B2)$  vs.  $\ln(B4) = 0.8147$ ; and  $\ln(B3)$  vs.  $\ln(B4) = 0.9089$ . Based on the processed data at the sand area at different depths, as illustrated in Fig. 6, and the use of Equations 4, 3, and 2, the variance, covariance, parameter 'a', and attenuation coefficient ratios values, and DII equations can be derived as presented in Table 5 below. Before creating the coral reef map using maximum likelihood classification, the next step after correcting the water column is masking. This process defines the boundaries of different objects in the image, such as separating the water body from the land to focus on the coral reef areas. In Fig. 7, the masking image shows that only the coral reefs have a digital reflectance value greater than zero, while the deep sea and land have zero values.

As the final stage of image processing, image classification consists of unsupervised and supervised classification (Fig. 8). The unsupervised classification was carried out to sort the visu-

**Table 4. The equation for the sun glint (de-glint) process**


Band	value slope (a)	Corrected de-glint equation
B2 (Blue)	1.1441	$Y = 6.9808 - 1.1441 \times (B5 - 1.4918)$
B3 (Green)	1.0145	$Y = 3.0609 - 1.0145 \times (B5 - 1.4918)$
B4 (Red)	1.0088	$Y = 1.1597 - 1.0088 \times (B5 - 1.4918)$



**Fig. 6. Plot bands pairs of ln (B3) vs ln (B2) (A), ln (B4) vs ln(B2) (B), and ln (B4) vs ln(B3) (C).**

**Table 5. Summarized values of the Variance, Covariance, “a” parameter, Attenuation coefficients, and DII equation generated from analysis of Landsat-8 OLI data**

Band	Variance	Band Pairs	Covariance	a	K <sub>i</sub> /K <sub>j</sub>	Depth in variant index (DII)
Ln (B2)	0.159	Ln (B2) vs Ln (B3)	0.108214	0.371	1.438	$Y = \ln(B2) - 1.438 \times \ln (B3)$
Ln (B3)	0.078	Ln (B2) vs Ln (B4)	0.071017	0.837	2.141	$Y = \ln(B2) - 2.141 \times \ln (B4)$
Ln (B4)	0.040	Ln (B3) vs Ln (B4)	0.052665	0.365	1.430	$Y = \ln(B3) - 1.430 \times \ln (B4)$

**Fig. 7. A mask image of the study sites shows coral reef areas that have a value > 0 (white color), while land and deep water have a value of zero (black color).**

al appearance of objects based on color hues into several classes on the water column-masked image. The iterative self-organizing for the unsupervised classification technique produces eight class groups (Fig. 8A), which can be used as a tentative map for field testing using 62 ground truth points. This image was then used as the input data to classify the habitat class in the water column corrected image (Fig. 7), and the output is a distribution map of five different habitat classes, algae, corals, seaweed, sand, and rubble (Fig. 8B).


Table 6 presents the results of simple calculations assessing the accuracy of five habitat classes based on data from 62 field checkpoints (ground truth). Meanwhile, Fig. 9 illustrates the accuracy test results using a confusion matrix calculation. The findings from the simple accuracy test of Table 6 indicate an

overall accuracy of 82.1%. The coral, seaweed, and sand classes have an accuracy of over 90%, with the coral class achieving the highest accuracy at 94.4%. In contrast, the algae and rubble classes display accuracies ranging from 62.5% to 66.7%, with the rubble class recording. The lowest accuracy was 62.5%.

The accuracy calculation from the confusion matrix (Fig. 9) shows an overall accuracy of 87.09% and a kappa coefficient of 0.83. Besides the overall accuracy, the confusion matrix calculation also includes two accuracy types: user accuracy, which indicates how often classified pixels are correct within their assigned class, and producer accuracy, which measures how well the classification aligns with the reference data (Congalton & Green, 2019). The user and producer accuracy values for benthic habitats vary except for sand. Producer accuracy is the highest for live coral (94.4%), followed by Seaweed (93.8%), Sand (92.9%), Algae (66.7%), and Rubble (62.5%). For user accuracy, Algae is the highest at 100%, followed by Sand (92.9%), Seaweed (88.2%), Rubble (83.3%), and Coral (81.0%). Based on the classification results, the overall accuracy of the coral reef habitat using 62 ground checkpoints was 87.1%. Thus, this study was reasonably accurate.

Table 7 presents the calculated areas of each benthic habitat at the study sites on the Kadatua, Liwutonkidi, and Siompu Islands based on the MLH classification of Landsat-8 OLI imagery, as illustrated in Fig. 8. From this table, coral is the most significant area, followed by sand, seaweed, rubble, and algae. In contrast, the island with the largest habitat is Kadatua, followed by Siompu, and the lowest is Liwutonkidi. The area of Kadatua Island is much smaller, only 37.9 km<sup>2</sup>, than that of Siompu Island, 112.8 km<sup>2</sup> (the Central Statistic Agency of South Buton Regency, 2024), while the coral cover of Kadatua, 173.2 ha, not that much different from Siompu around 186.2 ha. However, the rubble on Siompu Island (104.1 ha) is almost twice as high as that on Kadatua Island (56.7 ha). According to Haruddin et al. (2011), this high coverage of rubble on Siompu Island, particularly in BTNS-17 is directly proportional to the reports that many fishermen still catch the fish using explosives and potassi-






**Fig. 8. Image classifications using Landsat-8 (OLI).** (a) resulting from unsupervised iterative self-organizing (ISO) data; (b) resulting from supervised maximum likelihood.

**Table 6. Simple accuracy calculation based on 62 ground truth points**

No.	Habitat class	Ground truth points	Accurate points	Accuracy (%)
1	Coral	18	17	94.4
2	Seaweed	16	15	93.8
3	Rubble	8	5	62.5
4	Sand	14	13	92.9
5	Algae	6	4	66.7

**Table 7. The benthic habitat (ha) areas in Kadatua, Liwutonkidi, and Siompu Islands based on the maximum likelihood classification of Fig. 8**

Island	Algae	Corals	Seaweed	Sand	Rubble	Total
Kadatua	33.64	173.18	86.96	97.33	56.74	447.85
Liwutonkidi	6.03	84.35	20.38	18.72	13.57	143.05
Siompu	33.01	186.19	74.32	95.89	104.13	493.54
Total	72.68	443.72	181.66	211.94	174.44	1,084.44
Percentage (%)	6.7	40.92	16.75	19.54	16.09	100.00



**Fig. 9. The accuracy of the benthic habitat classification of the study sites based on matrix confusion calculation.**

um cyanide.

### Coral reef data analysis

Table 8 presents the data collected from the fieldwork based on underwater photo transects of benthic live forms. The Hard coral (HC), which is composed of *Acropora* (AC) and non-*Acropora* (NAC), is considered quite reasonable based on the criteria of Giyanto et al. (2017) in Table 1 at the study sites of BTNC-D, BTNC-C, and BTNC-13 on Siompu, Liwutonkidi, and Kadatua Islands that have a percentage of HC > 35%, but on the other study sites like BTN-17 and BTNC-14 were classified as moderate, ranging between 19 and 35%. The combination of FS and dead coral with algae in all study sites is actually in moderate

**Table 8. Benthic live form (%) collected using underwater photo transects based on LIT's field works**

Benthic life form	Siompu Island		Liwutonkidi Island	Kadatua Island		Average
	BTNC-D	BTNC-17	BTNC-C	BTNC-13	BTNC-14	
Hard coral (HC = AC + NAC)	43.8	23.5	44.5	40.1	33.5	37.1
<i>Acropora</i> (AC)	11.5	4.1	11.1	7.7	27.3	12.3
Non <i>Acropora</i> (NAC)	32.3	19.4	33.3	32.4	6.3	24.8
Recent dead coral (DC)	0.1	0.0	0.1	0.1	0.3	0.1
Dead coral with algae (DCA)	48.6	17.6	47.9	23.8	39.9	35.6
Soft coral (SC)	0.5	0.3	0.1	0.1	1.2	0.4
Sponge (SP)	3.4	6.0	3.2	2.7	11.1	5.3
Fleshy seaweed (FS)	0.0	0.0	0.0	0.0	0.0	0.0
Other biota (OT)	0.1	0.8	0.5	0.7	0.6	0.6
Rubble (R)	1.8	48.7	1.3	31.3	13.5	19.3
Sand (S)	1.7	3.1	2.5	1.1	0.0	1.7
Silt (SI)	0.0	0.0	0.0	0.1	0.0	0.0
Rock (RK)	0.0	0.0	0.0	0.0	0.0	0.0
Total	100.0	100.0	100.0	100.0	100.0	100.0

condition, but since the criteria of Table 2 are classified only into two classes of high and low, thus all moderate classes are classified as High.

The third criterion for the CHI is the density of coral reef biomass (kg/ha), as shown in Table 1. Thus, Table 9 lists the coral fish in all study sites. The coral reef fish consisted of two groups, the carnivorous and the herbivorous. The fish belongs to carnivorous include the families of Seranidae (Groupers), Lutjanidae (Snappers), Lethrinidae (Emperors or Scavengers), and Haemulidae (Grunts). Meanwhile, the herbivores are the families of Acanthuridae (Surgeonfishes, Tangs, and Unicornfishes), Scaridae (Parrot fish), and Siganidae (Rabbitfish). The total fish biomass density at the sampling locations from highest

to lowest in order is BTNC-13 (895.2 kg/ha), BTNC-D (722.8 kg/ha), BTNC-C (446.9 kg/ha), BTNC-17 (391.7 kg/ha) and the lowest BTNC-14 (329.3 kg/ha). Based on the criteria of the CHI in Table 1, all the study sites fall outside the category of coral with high or medium health index because the RFB at all study sites was much lower than 940 kg/ha, so it is classified as low. The low RFB in all study sites was due to the lousy exploitation of reef fish by unfriendly fishing methods, mainly explosives and poison (Potassium cyanide; Haruddin et al. 2011).

Table 10 summarizes the calculation of the CHI based on the benthic live forms data collected during the underwater photos transect (Fig. 3 and Table 8) and RFB of the carnivorous and herbivorous (Fig. 4 and Table 9). According to the criteria

**Table 9. Biomass (kg/ha) of coral reef fish belongs to the Carnivorous and Herbivorous groups**

No.	Coral Fish Family	Siompu Island		Liwutonkidi Island	Kadatua Island	
		BTNC-D	BTNC-17	BTNC-C	BTNC-13	BTNC-14
Carnivorous						
1	Seranidae	52.1	20.9	24.8	54.1	5.9
2	Lutjanidae	155.7	25.8	7.9	151.3	2.2
3	Lethrinidae	82.3	77.8	5.6	107.7	2.2
4	Haemulidae	36.1	6.9	2.6	30.1	11.2
Herbivorous						
5	Acanthuridae	157.4	181.6	301.1	313.5	278.5
6	Scaridae	172.1	69.2	83.9	190.0	11.7
7	Siganidae	67.1	9.5	21.0	48.4	17.6
Total biomass		722.8	391.7	446.9	895.2	329.3

**Table 10. The coral health index (CHI) of the study sites based on the percent live coral cover (LCC), flashy algae cover of coral resilience potency, and coral fish biomass (CFB)**


Study sites		Live benthic Forms Category				Resilience	Resilience		Coral health index (CHI)
		HC	FS	R	% Live coral cover (LCC)		Biomass (kg/ha)		
BTNC-D	Siompu Is.	43.8	−0	1.8	High	High	722.8	Low	6
BTNC-17	Siompu Is.	23.5	−0	48.7	Moderate	High	391.7	Low	5
BTNC-C	Liwutonkidi	44.5	−0	1.3	High	High	466.9	Low	6
BTNC-13	Kadatua Is.	40.1	−0	31.3	High	High	895.2	Low	6
BTNC-14	Kadatua Is.	33.5	−0	13.5	Moderate	High	329.3	Low	5

HC, hard coral; FS, fleshy seaweed; R, rubble.

of the CHI (Table 2), the CHI in the study sites ranges from value Index of 5 on the BTNC-17 (northern Siompu Island) and BTNC-14 (southern Kadatua Island) to Index value 6 on BTNC-D (southern of Siompu Island), BTNC-C (Liwutonkidi), and BTNC-13 (northern of Kadatua Island). The result of this analysis is mapped in Fig. 10. These CHI values of 5 were categorized as a little bit higher than poor, while 6 were classified as medium level, indicating that the coral reef was recovering. Hence, better management is needed to achieve better health conditions.

## Discussions

This study discusses the mapping of coral reef habitats in Kadatua, Liwutonkidi, and Siompu Islands using Landsat-8 OLI satellite imagery data and further applies it to develop a CHI assessment. The standard procedures for satellite image processing are applied, such as corrections for atmospheric, sun glint, and water column influences. Habitat classification by applying the conventional method was then done using MLH (Fig. 1). The classification map consists of 5 classes, Algae, Coral, Seaweeds, Sand, and Rubbles, with the area for each class can be calculated



**Fig. 10. Coral reef map.** (A) maximum likelihood Classification of benthic habitat based on Landsat-8 (OLI); (B) coral health index (CHI) based on benthic live form and coral reef fish abundance data obtained from field work using (LIT) method.

(Table 7 and Fig 8) and the accuracy of the classification map is also determined (Fig. 9).

Regarding map accuracy, the accuracy value will be higher if the agreement between the classification results and field observations is higher (Hossain et al., 2020). Therefore, selecting an inappropriate ROI on each class representative can reduce the producer accuracy value, which indicates that all habitats generated from the image can be mapped well. Also, the user accuracy values show that the results of field observation of habitat classes can be used correctly in the mapping process (Hafizt et al., 2017).

In this study, the results of the simple classification of the coral reef habitat using Landsat-8 OLI image showed an overall accuracy of 87.2% within 62 ground truth points used, while based on the confusion matrix, the overall accuracy is 87.1%. The lowest accuracy is the producer accuracy of 62.5% for Rubble. Meanwhile, the lowest user accuracy of 81.0% is for corals and the kappa coefficient of 0.83. Based on the rules contained in Indonesian National Standard (SNI) of 7716:2011 (LIPI, 2014), the acceptable accuracy value for shallow seabed habitat mapping is  $\geq 60\%$ . According to Mumby et al. (1998), a 65-70% accuracy for mapping aquatic habitats using satellite sensing can be reasonably good. Furthermore, a 60-80% accuracy can be used to recommend natural resource monitoring inventory activities (Green et al., 2000). Therefore, in this study, the results of this classification for coral reef mapping were accurate enough and able to be used for various management purposes. Regarding the condition of coral reef, such as percent of LCC based on the field observation using LIT method indicated that out of the five observation sites, four of them (BTNC-D/Kadatua Island, BTNC-C/ Liwutonkidi Island, BTNC-13 and -14/ Siompu Island) were in moderate condition (LCC in the ranges of 25–49.9%), except BTNC-17/Kadatua Island was in poor/bad condition (LCC < 25%). However, if the detailed analysis, such as Coral Mortality Index ( $CMI = \% \text{ dead coral} / \% [\text{Dead} + \text{Live corals}]$ ) and Coral Bio-Erosion Index ( $CBI = \% \text{ HC} / [\text{HC} + \text{soft coral} + \text{Sponge} + \text{algae} + \text{other}]$ ) were computed using data in Table 8, with the criteria of both CMI and CBI of 0 – < 0.25 is quantified as 4; 0.24–0.50 as 3; 0.50–0.75 as 2; and > 0.72 as 1, then the sum of both criteria, resulting the ecological status of Very poor coral condition if the value: 0–2, Poor: 3–4, Good: 5–6 and Excellent: 7–8 (English et al., 1999; Zamani & Januar, 2020). Based on the above criteria and the computation, all the study sites are in the ranges of value 3 (BTNC-D, -C, and -14) and value 4 (BTNC-17 and -13). This fact revealed that the cor-

al conditions in all observation sites are poor, which explains the tendency of HCs to shift into non-coral builder organisms, such as soft coral, sponges, algae, and others (Zamani & Januar, 2020). These poor coral conditions are strongly related to lousy fishing practices used by the fishermen, especially in BTNC-17 (Haruddin et al. 2011). Analysis of the CHI in Table 10 also showed that sites of BTNC-17 have the lowest CHI values of 5, followed by BTNC-14 of 5. These two sites also have the lowest RFB of < 400 kg/ha compared to the other observation sites.

Coral reefs are home to the world's resources, and millions of people depend on coral for their livelihoods. However, local stressors and climate change have forced coral reefs to live under pressure (Souter et al., 2021). This study shows that the local stressors that most cause coral reef damage are destructive fishing practices, such as the use of explosives, cyanide, and environmentally unfriendly fishing gear that destroy coral reef structures and reduce biodiversity; in addition, the practice of taking coral for construction materials, decoration purposes, and the aquarium trade, weakening the ecosystem (Haruddin, 2011; Fig. 11). Furthermore, global stressors to coral reefs are caused by coral bleaching. Although there are no specific reports of coral bleaching events on the islands of Kadatua, Liwutonkidi, and Siompu, in general, coral bleaching has been observed in areas around those islands, such as the Wakatobi Islands, especially during El Niño events, which increases sea surface temperatures (SST) in 2010 and 2016 (Wouthuyzen et al. 2018). So, the coral bleaching events on the Kadatua, Liwutonkidi, and Siompu Islands, which are part of an ecosystem close to and similar to the Wakatobi Islands, are likely to cause coral bleaching.

The explosives utilized to exploit coral reef fish can cause coral reef degradation characterized by low percentages of LCC, high dominance of rubble and algae, or even mass coral death (Fig. 11). Benthic algae will quickly colonize the dead coral reefs (Liao et al., 2021). This can be seen from the results of underwater photo transects in the benthic life form data of Table 8, where the combined components of rubble and dead coral covered with algae in almost all study locations are high, ranging from 50–60%. Still, at the study site in BTNC-17 (Siompu Island), it is the highest at 66.3%, as well as the cover of live coral reefs, is classified as poor (LCC is <25%). However, the other four locations are in moderate condition (25–50%).

Coral reef quality degradation is a painful truth, considering that the growth of coral is relatively slow (Fong & Todd, 2021; Morais et al., 2020), especially the *Porites heronensis*





**Fig. 11. Coral damage due to destructive fishing practices in study sites of BTNC-17.**

during the winter months ( $0.48 \pm 0.04$  mm per month) and *Pocillopora damicornis* during the summer months ( $1.18 \pm 0.08$  mm per month; Anderson et al., 2015). Corals that experience a decline in quality of life tend not to be able to provide ecosystem goods and services as they should. This study found that dynamite fishing to get fish is much more detrimental to coral reefs because reefs are home to resources that are needed by all living creatures. Damaged coral ecosystems are linear with low fish production, such as the condition in Fig. 11, which does not show any fish, even one. The high intensity of coral reef degradation in the long term will eliminate biodiversity, reduce fish catches, and ultimately place coastal welfare in an irreparable position, particularly for communities that depend on coral reef ecosystem services and marine resources (Benkwitt et al., 2020).

Reef fish coexist with coral reefs. Thus, the diversity and biomass of fish in coral reef ecosystems directly reflects the health of coral reefs. In this study, the biomass density of reef fish that consisted of 2 groups, the carnivorous (4 Families) and the Herbivorous (3 Families) (Table 9), was used to assess the coral reef's health by applying the criteria of fish biomass density of  $> 1,940$  kg/ha as the healthy coral reef environment, otherwise,  $< 940$  as a poor condition, while in between those values as a medium condition (Table 1; Giyanto et al., 2017). The entire reef fish in all study sites was below the criteria (Table 9), which means that judging by coral reef fish abundance only, all

coral reef environments in the study sites were in poor condition. Thus, the low coral fish below the criteria for healthy coral reefs due to dynamite and poison fishing that cause damage to benthic habitat and exacerbated by overfishing reveals facts that the coral conditions at the study sites are low health condition (Carvalho et al., 2021), such as at the BTNC-17 (Siompu Island) to moderate conditions for other study sites.

Table 11 lists the criteria for determining the health of coral reefs in the Caribbean waters of the Mesoamerican region (Mexico, Belize, Honduras, and Guatemala) based on the biomass of herbivorous and carnivorous coral fish (see the criteria in the notes below Table 11) as a comparison to the criteria developed by Giyanto et al (2017). Unlike the criteria of Giyanto et al (2017), in this criteria, the herbivorous fish used are limited to the surgeonfish group (family Acanthuridae, especially from the species *Acanthhurus* spp and *zebrasoma* spp) and the parrotfish group (Family Scaridae, such as *Cetoscarus* spp; *Chlorurus* spp; *Hipposcarus* spp; *Scarus* Spp.). These Herbivorous fish are essential grazers in controlling the macroalgae that could overgrow the reef. Meanwhile, the carnivorous fish group is limited to fish from the family Serranidae (groupers) and Lutjanidae (Snapper). According to Table 11, the condition of coral reef fish from the Herbivore group is classified as high for BTNC-D and -13 and poor for BTNC-17, -C, and -14, while the Carnivore group is grouped as very high for BTNC-D, and -13, Poor for BTNC-17,



**Table 11. Coral reef fish in study sites according to the criteria of Mesoamerican**

Coral reef-fish group	BTNC-D	BTNC-17	BTNC-C	BTNC-13	BTNC-14
Herbivores (all surgeonfish, and all Parrotfish):					
Acanthuridae ( <i>Acanthurus</i> spp; <i>Zebrazoma</i> spp)	5,327	2,160	2,934	4,859	3,071
Scaridae ( <i>Cetoscarus</i> spp; <i>Chlorurus</i> spp; <i>Hipposcarus</i> spp; <i>Scarus</i> Spp.)	6,025	2,423	2,937	6,650	408
Total Herbivore fish (g/350 m <sup>2</sup> )	11,352	4,583	5,871	11,509	3,479
Total Herbivore fish (g/100 m <sup>2</sup> )	3,243	1,309	1,677	3,288	994
Herbivore coral reef fish condition	Good	Poor	Poor	Good	Poor
Carnivores (all families of Serranidae and Lutjanidae):					
Serranidae (Grouper)	52,114	731	867	1,895	207
Lutjanidae (Snapper)	5,448	904	277	5,297	78
Total Carnivore fish (g/350 m <sup>2</sup> )	57,562	1,635	1,144	7,192	285
Total Carnivores fish (g/100 m <sup>2</sup> )	16,446	467	327	2,055	81
Carnivore coral reef fish condition	Very good	Poor	Critical	Very Good	Critical

Criteria for Herbivores and Carnivores.

Herbivore fish (g/m<sup>2</sup>): Critical, <990; Poor, 990–< 1,860; Fair, 1,860–< 2,740; Good, 2,740–< 3,290; Very good, ≥ 3,290.

Carnivore fish (gram/m<sup>2</sup>): Critical, <390; Poor, 390–<800; Fair, 800–< 1,210; Good, 1,210–< 1,620; Very good, ≥ 1,620.

Data from McField et al., (2024)

and critical for BTNC-C and -14. Thus, the result of CHI determination based on Mesoamerican criteria, in general, is almost the same as Giyanto's et al (2017) criteria for BTNC-17, -C, and -14 but the criteria for the Mesoamerican region seem more manageable to analyze, more detailed and reliable, and simpler than the criteria developed by Giyanto et al (2017), since it use only data of 2 groups of herbivore (Acanthuridae and Scaridae), and 2 groups carnivore fishes (Lutjanidae and Serranidae).

In this study, it was seen that CHI was highly correlated with reef fish abundance. For example, although the CHI level in BTNC-D, -C, and -13 was 6, while BTNC-17 and BTNC-14 were 5, all study sites had a low fish abundance of < 970 kg/ha (Table 9) based on the criteria of Giyanto et al. (2017; Table 10). These results of CHI computation would be different if the fish abundance criteria from McField et al. (2024), were used (Table 11), which showed that BTNC-D and -13 had fish abundance classified as “Good for the Herbivorous fish group” and “Very good for the Carnivorous fish group”, while the abundance of coral fish in BTNC-17, -C and -14 were classified as “poor” and “Critical” (Table 11). So, suppose the data in Tables 8 and 9 are reanalyzed using the criteria of McField et al. (2024); then, the analysis results reveal a more apparent difference in CHI of the two study sites (BTNC-D and -13), which have coral fish abundance in the categories “very good” and “good”, where the CHI in both locations is also classified as “very good” (Table 12). In contrast, the remaining 3 study sites (BTNC-17, -C and -14) with coral fish abundance in the categories “poor and Critical”

have CHI values classified as “fair/moderate (Table 12).

From the results of this analysis, it can be considered that the McField et al. (2024) criteria, such as in Table 12, are more reliable to be applied in determining the CHI in each study site and supporting the explanation of reef fish abundance than the Giyanto et al. (2017) criteria in Table 2, which classifies CHI with a wide range level from 10 to 1. Level 10 is the healthiest coral reef, with a high percentage of LCC and coral resilience, thus supporting abundant reef fish. Conversely, CHI level 1 is a coral reef with the lowest LCC and resilience, making it difficult for corals to grow and develop, resulting in a low abundance of economically valuable reef fish. Furthermore, there is no specific information about CHI for levels outside of the ranges from 10 to 1, for example, levels 9 to 2. So, it isn't easy to get detailed information on the status of the CHI.

The quantity and quality of LCC illustrate the level of resilience and affect how the benthic components survive. Corals with high levels of resilience have good adaptability when faced with disturbances and pressures. From Table 10, it can be seen that the coral resilience in the study sites is high, as indicated by the low fleshy algae cover (– 0%). However, the coral RFB based on Giyanto's et al (2017) criteria was low, while LCC varies from poor conditions, such as in BTNC-17, to moderate in other locations. The results of this study produce relatively a low CHI of 5 (BTNC-17 and -14), but a slightly high of 6 (BTNC-D, -C and -13, Table 13). Table 13 also shows that the CHI in this study was lower compared to the CHI of 2016 as in BTNC-14

**Table 12. Result of coral health index (CHI) reanalysis of data in Table 8 and 9 (lower panel) using the criteria of McField et al. (2024) (upper panel)**

CHI Grade / Level	CHI grade / Level ranges	LCC (%)	FMC (%)	HFB (g/m <sup>2</sup> )	CFB (g/m <sup>2</sup> )
Very good/excellent <sup>1)</sup>	4.3–5.0	> 40: (5)	< 2: (5)	> 3,290: (5)	> 1,620: (5)
Good <sup>2)</sup>	3.5–4.2	20–40: (4)	2–5: (4)	2,740–3,290: (4)	1,210–1,620: (4)
Fair <sup>3)</sup>	2.7–3.5	10–20: (3)	5–12: (3)	1,860–2,740: (3)	800–1,210: (3)
Poor <sup>4)</sup>	1.9–2.6	5–10: (2)	12–25: (2)	990–1,860: (2)	390–800: (2)
Critical <sup>5)</sup>	1.0–2.8	< 5: (1)	> 25: (1)	< 990: (1)	< 390: (1)
Study Sites	Average grade for CHI*	LCC Grade	FMC Grade	HFB Grade	CFB Grade
BTNC-D (Siompu Is) <sup>1)</sup>	4.75; Very good <sup>1)</sup>	43.8 (5) <sup>1)</sup>	–0: (5) <sup>1)</sup>	3,243 (4) <sup>2)</sup>	16,466 (5) <sup>1)</sup>
BTNC-13 (Kadatua Is) <sup>1)</sup>	4.75; Very good <sup>1)</sup>	40.1 (5) <sup>1)</sup>	–0: (5) <sup>1)</sup>	3,288 (4) <sup>2)</sup>	2,055 (5) <sup>1)</sup>
BTNC-17 (Siompu Is) <sup>3)</sup>	3.25; Fair <sup>3)</sup>	23.5 (4) <sup>2)</sup>	–0: (5) <sup>1)</sup>	1,309 (2) <sup>4)</sup>	467 (2) <sup>4)</sup>
BTNC-C (Liwutonkidi Is) <sup>3)</sup>	3.25; Fair <sup>3)</sup>	44.5 (5) <sup>1)</sup>	–0: (5) <sup>1)</sup>	1,677 (2) <sup>4)</sup>	327 (1) <sup>5)</sup>
BTNC-14 (Kadatua Is) <sup>3)</sup>	3.0; Fair <sup>3)</sup>	35.5 (4) <sup>2)</sup>	–0: (5) <sup>1)</sup>	994 (2) <sup>4)</sup>	81 (1) <sup>5)</sup>

The number in the bracket denotes the grade (Very good, Good, Fair, Poor, Critical) of the LCC, FMC, HFB, CFB.

The average grade of CHI = (Grade of LCC + Grade of FMC + Grade of HFB + Grade of CFB) / 4.

\* The average grade of CHI for BTNC-D = (5 + 5 + 4 + 5) / 4 = 4.75 (Very good).

<sup>1)</sup> Very good (excellent).

<sup>2)</sup> Good.

<sup>3)</sup> Fair.

<sup>4)</sup> Poor.

<sup>5)</sup> Critical.

LCC, live coral cover; FMC, Fleshy macro-algae cover; HFB, herbivore fish biomass; CFB, carnivore fish biomass.

**Table 13. Comparison of the conditions of coral health index (CHI) in the study sites between 2016 and 2021**

Study sites		Coordinates		Coral health index (CHI)	
		Latitude	Longitude	2016*	2021**
BTNC-D	Siompu Island	–5.67623	122.46689	5	6
BTNC-17	Siompu Island	–5.62051	122.51731	5	5
BTNC-C	Liwutonkidi Island	–5.60271	122.49907	6	6
BTNC-13	Kadatua Island	–5.51706	122.47507	6	6
BTNC-14	Kadatua Island	–5.56261	122.48058	6	5

\* Data from Giyanto et al. (2017).

\*\* This study.

from a level of 6 to 5. However, there were also increases, such as BTNC-D from a level index of 5 to 6, while the remaining BTNC-C and -13 had the same level of 6 both in 2016 and 2021 including BTNC-17 that surprisingly remain the same as in 2016. Therefore, the CHI on the study sites, in general, tends to decrease, but some exceptions show also improvement, such as in BTNC-D or remain the same as 2016. Despite high coral resilience indicated by low fleshy algae presence (resilience potency), the overall CHI remains impacted by low coral RFB and variable of coral cover conditions.

To address this decline, particularly for the study sites, several efforts can be made as follows: (1) To control lousy fishing practices and overfishing in these study sites by implementing

and enforcing sustainable fishing to maintain the balance of marine life; (2) To reduce marine pollution by limiting the discharge of pollutants, including sewage, agricultural runoff, and sedimentation into the ocean. (3) To allow coral reefs to recover and thrive without human interference, e.q. by establishing marine protected areas (MPA) or No-Take Zone areas; (4). To Engage in coral restoration projects, such as coral gardening and transplantation, to help damaged reefs recover; (5). To educate and involve the local communities in conservation efforts to ensure long-term sustainability and protection of coral reefs. These efforts can help improve the CHI and ensure the long-term health and resilience of coral reef ecosystems.

Special efforts must be emphasized to rehabilitate the

coral reef degradation, especially in the three sites, the BTNC-17, -14, and -C, followed with the kind effort from the above recommendation of points 1, 2, and 5, but firstly by starting the establishment of the No-take zone or MPA (point 3) and conducting coral transplantation (point 4). The No-take zone/MPA established by the local community has proven effective in recovering both the reef condition and reef fish population in the 20 sites of the small Islands of the Biak-Numfor and Supiori Districts in the Gulf of Cendrawasih, Papua in 2009–2014. The abundance of reef fish inside the No-take zone was 3–4, 3–5, and 2–3 times higher than the outside for the target/commercial fishes: the fish being the catch target by fishers, indicator fish, the butterfly fish of the family Chaetodontidae, which can indicate the health of the coral reefs; and major fish group, the relatively small size fish that are mostly grouped as ornamental/aquarium fishes, respectively (Wouthuyzen et al. 2016). The two study sites that show good/excellent CHI, like BTNC-D and -13 (Table 12), are also better if the No-take zone is established there. It is hoped that the reef fish from the No-take zones can spill over to the surrounding sites or outside the established no-take zone.

## Conclusions

This study assesses the CHI using Landsat-8 OLI remote sensing and in-field LIT surveys. The Landsat-8 data effectively mapped benthic habitats, including coral, macroalgae, sand, and rubble, with – 81% overall accuracy. The LIT surveys provided data on LCC, fleshy macroalgae as a resilience indicator, and reef fish abundance (using UVC, Fig. 4), which were used to determine the CHI.

Based on the Giyanto et al. (2017) criteria, CHI was categorized into two levels: level 5 (BTNC-14 and -17) and level 6 (BTNC-13, -C, and -D). A comparison between 2016 and 2021 revealed a decline in CHI for BTNC-14 (level 6 to 5) and an improvement for BTNC-D (level 5 to 6), while other sites remained at level 6. However, distinguishing whether levels 5 and 6 indicate good or poor conditions remains challenging. Reef fish abundance was low (<970 kg/ha) across all sites, with BTNC-14, -D, and -C having the lowest values (<500 kg/ha), which is related to destructive fishing practices.

CHI was also compared using McField et al. (2024) criteria applied for annually monitoring the coral reefs in Mesoamerican (Mexico, Belize, Guatemala, and Honduras) countries from 2006 to 2023. This method classified BTNC-13 and -D as “very

good/excellent” and BTNC-C, -17, and -14 as “fair.” McField’s criteria provide a more precise, reliable framework that is simpler for assessing CHI and explaining reef fish abundance, making them a viable alternative for monitoring Indonesia’s coral reef ecosystems.

Based on these findings, several key recommendations are issued: (1) Establish a MPA by designating a small no-take zone of about 1 to 10 ha at each study site to protect the coral reefs, and allow the fish population to recover, and enhance fish spill-over into surrounding areas (this approach has proven successful in Biak Island, Papua; Wouthuyzen et al. 2016). (2). Implement coral transplantation/coral gardening, particularly in sites with poor or fair CHI levels, such as BTNC-14, -17, and -C. (3). Regulate fishing practices by enforcing sustainable fishing regulations to curb overfishing and destructive methods like blast and cyanide fishing, and ensure marine life balance. (4). Reduce marine pollution by minimizing the discharge of pollutants, including sewage, agricultural runoff, and sedimentation, to improve water quality and support reef resilience. (5). Enhance community involvement and monitoring by conducting long-term research and involving local communities in conservation initiatives to promote awareness and sustainable reef management.

Those recommendations provide a roadmap for improving reef health and ensuring the long-term sustainability of coral ecosystems in the Kadatua, Liwutonkidi, and Siompu Islands. Therefore, this study’s findings can serve as a benchmark for monitoring future changes in the coral reef environment.

## Competing interests

No potential conflict of interest relevant to this article was reported.

## Funding sources

We received financial support for this study from the Indonesian Institute of Sciences (LIPI) COREMAP-CTI 2021–2022, grant No. 17/A/DK/2021.

## Acknowledgements

We thank those who assisted us during the research, both in the field and beyond, though we cannot mention everyone. We also thank the anonymous reviewers and the FAS journal editor for their comments that improved this paper.

## Availability of data and materials

Upon reasonable request, the datasets used in this study can be made available from the corresponding author.

### Ethics approval and consent to participate

Not applicable.

### ORCID

Daniel Deonisius Pelasula

<https://orcid.org/0000-0002-6692-1510>

Sam Wouthuyzen <https://orcid.org/0000-0001-9655-6254>

Mufti Petala Patria <https://orcid.org/0000-0001-8648-9529>

Simon Izaak Patty <https://orcid.org/0009-0000-6270-2866>

Petrus Christianus Makatipu  
<https://orcid.org/0009-0004-2650-7122>

Baru Sadarun <https://orcid.org/0009-0001-6868-5128>

Wiesye Violent Pelupessy  
<https://orcid.org/0009-0005-8035-8380>

Suiyuan Wang <https://orcid.org/0000-0003-0720-8173>

Johanis Dominggus Lekallete  
<https://orcid.org/0000-0001-6280-7772>

## References

- Adji AS. Suitability analysis of multispectral satellite sensors for mapping coral reefs in Indonesia case study: Wakatobi Marine National Park. *Mar Res Indonesia*. 2014;39:73-8.
- Anderson KD, Heron SF, Pratchett MS. Species-specific declines in the linear extension of branching corals at a subtropical reef, Lord Howe Island. *Coral Reefs*. 2015;34:479-90.
- Aulia ZS, Ahmad TT, Ayustina TT, Hastono FT, Hidayat RR, Mustakin H, et al. Shallow water seabed profile changes in 2016-2018 based on Landsat 8 satellite imagery (case study: Semak Daun Island, Karya Island and Gosong Balik Layar). *Omni-Akuatika*. 2020;16:26-32.
- Benkwitt CE, Wilson SK, Graham NAJ. Biodiversity increases ecosystem functions despite multiple stressors on coral reefs. *Nat Ecol Evol*. 2020;4:919-26.
- Brodie JE, McElroy AE. Crown-of-thorns starfish and their role in coral reef degradation: a review. *Mar Ecol Prog Ser*. 2015;529:101-13.
- Burke L, Reyttar K, Spalding M, Perry A. Reefs at risk revisited in the coral triangle. Washington, DC: World Resources Institute (WRI); 2012.
- Carvalho PG, Setiawan F, Fahlevy K, Subhan B, Madduppa H, Zhu G, et al. Fishing and habitat conditions differentially affect the size spectra slopes of coral reef fishes. *Ecol Appl*. 2021;31:e02345.
- Chen A, Ma Y, Zhang J. Partition satellite-derived bathymetry for coral reefs based on spatial residual information. *Int J Remote Sens*. 2021;42:2807-26.
- Congalton RG, Green K. Assessing the accuracy of remotely sensed data. In: Principles and practices, 3rd ed. Boca Raton, FL: CRC Press; 2019.
- Cooper EL, Hirabayashi K, Strychar KB, Sammarco PW. Corals and their potential applications to integrative medicine. *Evid Based Complement Alternat Med*. 2014;2014:184959.
- Darling ES, D'agata S. Coral reefs: fishing for sustainability. *Curr Biol*. 2017;27:R65-8.
- English S, Wilkinson C, Baker V. Survey manual for tropical marine resources. 2nd ed. Townsville: Australian Institute of Marine Science; 1999.
- Fabricius KE. Effects of terrestrial runoff on the ecology of corals and coral reefs: review and synthesis. *Mar Pollut Bull*. 2005;50:125-46.
- Fell F. A contrast minimization approach to remove Sun glint in Landsat 8 imagery. *Remote Sens*. 2022;14:4643.
- Ferrario F, Beck MW, Storlazzi CD, Micheli F, Shepard CC, Airoidi L. The effectiveness of coral reefs for coastal hazard risk reduction and adaptation. *Nat Commun*. 2014;5:3794.
- Fong J, Todd PA. Spatio-temporal dynamics of coral-macroalgal interactions and their impacts on coral growth on urbanized reefs. *Mar Pollut Bull*. 2021;172:112849.
- Fox HE, Harris JL, Darling ES, Ahmadiya GN, Estradivari, Razak TB. Rebuilding coral reefs: success (and failure) 16 years after low-cost, low-tech restoration. *Restor Ecol*. 2019;27:862-9.
- Froese R, Pauly D. FishBase 2000: concepts, designs, and data sources. Penang: WorldFish; 2000.
- Giyanto, Soedarma D. Efisiensi dan akurasi pada proses analisis foto bawah air untuk menilai kondisi terumbu karang. *Oseanologi dan Limnologi di Indonesia*. 2010;36:111-30.
- Giyanto, Mumby P, Dhewani N, Abrar M, Iswari MY. Indeks Kesehatan Terumbu Karang Indonesia. Jakarta: LIPI Press; 2017. p. 9.
- Green EP, Mumby PJ, Edwards AJ, Clark CD. Remote sensing handbook for tropical coastal management. Paris: UNESCO; 2000.
- Hafizt M, Iswari MY, dan Prayudha B. Assessment of landsat-8

- classification method for benthic habitat mapping in Padaido Islands, Papua. *Oceanologi dan Limnologi di Indonesia*. 2017;2:1-13.
- Haruddin A, Purwanto E, Budiastuti MS, Si M. Dampak kerusakan ekosistem terumbu karang terhadap hasil penangkapan ikan oleh nelayan secara tradisional di Pulau Siompu Kabupaten Buton Propinsi Sulawesi Tenggara. *J Ekosains*. 2011;3:29-41.
- Haya LOMY, Fujii M. Mapping the change of coral reefs using remote sensing and in situ measurements: a case study in Pangkajene and Kepulauan Regency, Spermonde Archipelago, Indonesia. *J Oceanogr*. 2017;73:623-645.
- Hedley JD, Harborne AR, Mumby PJ. Technical note: simple and robust removal of sun glint for mapping shallow-water benthos. *Int J Remote Sens*. 2005;26:2107-12.
- Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, et al. Coral reefs under rapid climate change and ocean acidification. *Science*. 2007;318:1737-42.
- Hoeksema BW. (2007). Delineation of the Indo-Malayan centre of maximum marine biodiversity: The Coral Triangle. In: Renema W, editor. *Biogeography, time and place: distributions, barriers and islands*. Dordrecht: Springer; 2007. p. 117–178.
- Hossain MS, Muslim AM, Nadzri MI, Teruhisa K, David D, Khalil I, et al. Can ensemble techniques improve coral reef habitat classification accuracy using multispectral data? *Geocarto Int*. 2020;35:1214-32.
- Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, Baird AH, et al. Global warming and recurrent mass bleaching of corals. *Nature*. 2017;543:373-7.
- International Coral Reef Initiative (ICRI). ICRI Annual Report 2020: advancing the protection and conservation of coral reefs globally. International Coral Reef Initiative (ICRI); 2020.
- Jackson JBC, Kirby MX, Berger WH, Bjorndal KA, Botsford LW, Bourque BJ, et al. Historical overfishing and the recent collapse of coastal ecosystems. *Science*. 2001;293:629-37.
- Kartikasari A, Pristianto T, Hanintyo R, Ampou EE, Wibawa TA, Borneo BB. Representative benthic habitat mapping on Lovina coral reefs in Northern Bali, Indonesia. *Biodiversitas*. 2021;22:4766-74.
- Kiel S, Klemens J. *Drupella* spp. (Mollusca: Gastropoda) On Indonesian coral reefs: the effects of grazing on coral health. *Coral Reefs*. 2006;25:177-80.
- Kohler KE, Gill SM. Coral point count with Excel extensions (CPCe): a visual basic program for the determination of coral and substrate coverage using random point count methodology. *Comput Geosci*. 2006;32:1259-69.
- Kulbicki M, Guillemot N, Amand M. A general approach to length-weight relationships for New Caledonian fishes. *Cy-bium*. 2005;29:235-52.
- Lembaga Ilmu Pengetahuan Indonesia (LIPI). Panduan teknis pemetaan habitat dasar perairan laut dangkal. Coremap. Jakarta: P2O-LIPI; 2014.
- Liao Z, Yu K, Chen B, Huang X, Qin Z, Yu X. Spatial distribution of benthic algae in the South China Sea: responses to gradually changing environmental factors and ecological impacts on coral communities. *Divers Distrib*. 2021;27:929-43.
- Lyzenga DR. Remote sensing of bottom reflectance and water attenuation parameters in shallow water using aircraft and Landsat data. *Int J Remote Sens*. 1981;2:71-82.
- McField M, Soto M, Martinez R, Giró A, Guerrero C, Rueda M, et al. 2024 Mesoamerican reef report card. Healthy Reefs for Healthy People. 2024 [cited 2025 Feb 4]. <https://www.healthyreefs.org/en/healthy-reefs-data/report-cards>
- Moberg F, Folke C. Ecological goods and services of coral reef ecosystems. *Ecol Econ*. 1999;29:215-33.
- Morais RA, Depczynski M, Fulton C, Marnane M, Narvaez P, Huertas V, et al. Severe coral loss shifts energetic dynamics on a coral reef. *Funct Ecol*. 2020;34:1507-18.
- Mumby PJ, Green EP, Clark CD, Edwards AJ. Digital analysis of multispectral airborne imagery of coral reefs. *Coral Reefs*. 1998;17:59-69.
- Nurdin N, Komatsu T, Agus, Akbar MAS, Djalil AR, Amri K. Multisensor and multitemporal data from Landsat images to detect damage to coral reefs, small islands in the Spermonde Archipelago, Indonesia. *Ocean Sci. J*. 2015;50:317-325.
- Pandolfi JM, Bradbury RH, Sala E, Hughes TP, Bjorndal KA, Cooke RG, et al. Global trajectories of the long-term decline of coral reef ecosystems. *Science*. 2003;301:955-8.
- Rolim FA, Rodrigues PFC, Langlois T, Neves LM, Gadig OBF. A comparison of stereo-videos and visual census methods for assessing subtropical rocky reef fish assemblage. *Environ Biol Fishes*. 2022;105:413-29.
- Song C, Woodcock CE, Seto KC, Lenney MP, Macomber SA. Classification and change detection using Landsat TM data: When and how to correct atmospheric effects? *Remote Sens Environ*. 2001;75:230-44.



- Souter D, Planes S, Wicquart J, Logan M, Obura D, Staub F. Status of coral reefs of the World: 2020. Townsville: Global Coral Reef Monitoring Network (GCRMN) & International Coral Reef Initiative (ICRI); 2021.
- Spalding M, Ravilious C, Green EP. World Atlas of coral reefs. Berkeley, CA: University of California Press; 2001.
- Spalding M, Burke L, Wood SA, Ashpole J, Hutchison J, zu Ermgassen P. Mapping the global value and distribution of coral reef tourism. *Mar Policy*. 2017;82:104-13.
- Sutherland KP, Porter JW, Torres C. Disease and immunity in Caribbean and Indo-Pacific zooxanthellate corals. *Mar Ecol Prog Ser*. 2004;266:273-302
- Syakti Ad, Jaya JV, Rahman A, Hidayati NV, Raza'i TS, Idris F, et al. Bleaching and necrosis of staghorn coral (*Acropora formosa*) in laboratory assays: immediate impact of LDPE microplastics. *Chemosphere*. 2019;228:528-35.
- The Central Statistic Agency of South Buton Regency. Regional Statistics of Buton Regency 2024. Sulawesi Tenggara: Badan Pusat Statistik Kabupaten Buton; 2024.
- Wouthuyzen S, Abrar M, Corvianawatie C, Salatalohi A, Kusumo S, Yanuar Y, et al. The potency of Sentinel-2 satellite for monitoring during and after coral bleaching events of 2016 in some islands of Marine Recreation Park (TWP) of Pieh, West Sumatra. *IOP Conf Ser Earth Environ Sci*. 2019;284:012028
- Wouthuyzen S, Abrar M, Lorwens Y. A comparison between the 2010 and 2016 El-Niño-induced coral bleaching in the Indonesian waters. *IOP Conf Ser Earth Environ Sci*. 2018;118:012051.
- Wouthuyzen S, Lorwens Y, dan Hukom FD. Do Fish sanctuary effective to conserve reef fishes? A case study in Biak-Numfor and Supiori Districts, Papua. *J Lit Perikan Ind*. 2016;22:271-84.
- Wulandari P, Sainal, Cholifatullah F, Janwar Z, Nasruddin, Setia TM, et al. The health status of coral reef ecosystem in Taka Bonerate, Kepulauan Selayar Biosphere Reserve, Indonesia. *Biodiversitas*. 2022;23:721-32.
- Zamani NP, Januar HI. Coral mortality and bio-erosion index for assessing environmental stress effects: a study case of the Indonesian tropical reef in Banda-Neira Conservation Park. *AACL Bioflux*. 2020;13:1027-1037.